

Which treatment protocol, among classical methods and/or various laser applications is the most effective in root canal disinfection, *in vitro*-? A systematic review

Joanna Theodosopoulou, Konstantinos Chochlidakis, Alexandra Tsigarida

¹Institute of Systems Biology, Shanghai University, Shanghai, China;
Email: joanna_theodosopoulou@yahoo.gr

Received ***** 2011.

ABSTRACT

Purpose: The aim of this systematic review was to answer the question “Which treatment protocol, among classical methods and/or various laser applications is the most effective in root canal disinfection, *in vitro*”. **Materials and Methods:** A MEDLINE, a Cochrane and an Embase search (three specified searches) were conducted to identify randomized controlled trials (RCT) until June 2010, conducted on human teeth and published in English, German or French language, examining the root canal disinfection after the use of lasers with or without mechanical instrumentation. Additionally, hand search was conducted and contact with authors, when needed. **Results:** The MEDLINE, the Cochrane and the EMBASE search identified 240, 28, and 35 published articles, respectively. Ten articles from the MEDLINE and 5 articles from the Cochrane search (that were also identified in the MEDLINE search) met the inclusion and validity assessment criteria. In *E.faecalis* elimination, instrumentation of the root canal and diode laser / 665nanometer / 1Watt (diode laser / 665nm / 1W) ir- radiation with the combined effect of Methylene Blue (MB) as photosensitizing agent (logCFU / ml = 1,636) seemed to be the best method. In *P.aeruginosa* and in *A.naeslundii* elimination, instrumentation of the root canal followed by irrigation with 5, 25%NaOCl (logCFU / ml = 0) seemed to be the best method. In general, instrumentation of the root canal followed by irrigation with 5, 25%NaOCl (logCFU / ml = 0) and instrumentation of the root canal and Er: YAG laser / 2940 nm / 0,8W irradiation (logCFU / ml = 1,924) seemed to be the best (polymicrobial studies). **Conclusion:** There are treatment protocols with the assistance or not of laser irradiation that can eliminate *E.faecalis*, *E.coli* and *S.aureus* inside the root canal. How-

ever, there is a serious number of *S. anginosus*, *F. nucleatum*, *A. naeslundii* and *P.aeruginosa* that remain inside the root canal even after laser irradiation. New research is needed in order to set a treatment protocol effective in the root canal disinfection from all bacteria that are related to endodontic origin pathology.

Keywords: Lasers; Classical Methods; Endodontic; Root Canal Therapy; Dentin; Bacteria; Disinfection

1. INTRODUCTION

One of the most crucial and fundamental stages of endodontic therapy is the root canal disinfection in its three-dimensional network of dentinal tubules. Nevertheless, persistence of infection in the root canal is responsible for long-term failures and need for endodontic therapy retreatments.

It is generally accepted that microorganisms tend to remain in the root canal even after proper preparation and are responsible for flare-ups, after the completion of the endodontic therapy. The most common of these microorganisms are: *Fusobacterium nucleatum*, *Enterococcus faecalis*, *Prevotella intermedia*, *Streptococcus anginosus*, *Treponema denticola*, *Porphyromonas gingivalis* (1-4).

During the last years, laser irradiation has been additionally introduced in root canal preparation, trying to gain acceptance for its disinfection ability in comparison with the common mechanical instrumentation and irrigation procedures.

Many studies examine the effectiveness of Nd: YAG, diode, Er,Cr:YSGG and Er:YAG laser, -when used in different wavelengths, solely, or in addition with various solutions- in the bacterial elimination inside the root canals.

The purpose of this systematic review was to answer the question “Which treatment protocol, among classical methods and / or various laser applications is the most ef-

fective in root canal disinfection, *in vitro*".

2. MATERIALS AND METHODS

2.1. Literature Search

One electronic search of MEDLINE from 1966 to June 2010 (错误! 未找到引用源。), one Cochrane (**Table 2**) and one Embase search from 1945 to June 2010 (**Table 3**) were conducted.

Table 1. Medline search strategy.

#	Search history	Results
#13	Search #12 Limits: Humans, English, French, German, Greek, Modern	240
#12	Search #11 AND #1	282
#12	Search #11 AND #1	13136
#11	Search #7 AND #10	15641
#10	Search #8 OR #9	14900
#9	Search (root canal therapy)	1622
#8	Search (endodontically treated teeth)	4830685
#7	Search #1 OR #2 OR #3 OR #4 OR #5 OR #6	138039
#6	Search (tooth OR teeth)	1267279
#5	Search bacteria	3639649
#4	Search method	19546
#3	Search dentin	14477
#2	Search endodont*	137179
#1	Search laser	240

Table 2. Cochrane search strategy.

#	Search history	Results
#1	laser	6294
#2	endodont*	1174
#3	dentin	1427
#4	method	241235
#5	bacteria	4978
#6	(tooth OR teeth)	7062
#7	(#1 OR #2 OR #3 OR #4 OR #5 OR #6)	248592
#8	(endodontically treated teeth)	89
#9	(root canal therapy)	401
#10	(#8 OR #9)	442
#11	(#7 AND #10)	422

#12	(#1 AND #11)	28
-----	--------------	----

Table 3. Embase search strategy.

#	Search history	Results
1	laser	106390
2	endodont*	1076
3	dentin	2983
4	method	785208
5	bacteria	121213
6	(tooth OR teeth)	40114
7	(#1 OR #2 OR #3 OR #4 OR #5 OR #6)	5138833
8	(endodontically treated teeth)	43
9	(root canal therapy)	64
10	(#8 OR #9)	1426192
11	(#7 AND #10)	218687
12	(#1 AND #11)	225630
13	limit 12 to humans, English or French or German or Greek from "1945 - 2007"	35

2.2. Inclusion Criteria-Validity

Three independent reviewers examined all the identified abstracts to determine whether they met the following criteria:

- 1) Study *in vitro*.
- 2) Conducted in human teeth.
- 3) Related to the question.
- 4) Experimental and control group.
- 5) Quantitative results provided.
- 6) English, German, French languages.

Whenever it was not possible to make this determination, the article was examined in full text. Subsequently, all relevant articles were obtained and a determination whether or not they met the inclusion criteria was made by three reviewers. It is important to state that only studies measuring the bactericidal effect of lasers and other procedures were included. The studies that examined the removal of smear layer or debris, the morphological or histological changes, the apical leakage after obturation and the dentin permeability were excluded, as irrelevant to our question (**Tables 8-12**).

All articles that met the inclusion criteria were assessed for validity. Validity was determined on a 7-point scale (**Table 4**) and studies not meeting 5 or more of the 7 validity criteria were excluded.

All articles were classified by evidence level (**Table 5**) (EBM://cebm.jr2.ox.uk/docs/levels.html) and then assessed

sed for Validity (<http://www.cebm.utoronto.ca/teach/materials/therapy.htm>) (Table 6).

Table 4. Validity assessment criteria¹.

1. Was the assignment of patients of treatment randomised?
2. Was the randomisation list concealed?
3. Was the follow-up of patients sufficiently long and complete?
4. Were all patients analysed in the groups to which they were randomised?
5. Were patients and clinicians blinded to the treatment being received?
6. Aside from the experimental treatment, were the groups treated equally?
7. Were the groups similar at the start of the trial?

Table 5. Clinical evidence levels².

Level of evidence	Study Type	#
1A	Randomized control trial (RCT) Systematic Review of RCTs	6
1B	Controlled Trial Systematic review of CCTs	2
2	Cohort Study(CCS) Systematic review of CSs	2
3	Case Control Study Systematic Review of CCSs	0
4	Case Series	0
5	Expert's Opinion Narrative Review	0
NA ³	Cross Sectional Case Reports Animal Studies Laboratory Studies	0
Non validated	Because of language limitations	0

Table 6. Validity assessment criteria application.

Reference	1	2	3	4	5	6	7
De Souza EB. <i>et al.</i> , 2008	Y	Y	Y	Y	N	Y	Y
Fimple JL. <i>et al.</i> , 2008	Y	Y	Y	Y	N	Y	Y
Fonseca MB. <i>et al.</i> , 2008	N	N	Y	Y	N	Y	Y
Bergmans L. <i>et al.</i> , 2008 ⁴	Y	Y	Y	Y	N	Y	Y
Foschi F. <i>et al.</i> , 2007	Y	Y	Y	Y	N	Y	Y
Wang QQ. <i>et al.</i> , 2007 <small>错误! 未定义书签。</small>	Y	Y	Y	Y	N	Y	Y
Gordon W. <i>et al.</i> , 2007	N	N	Y	Y	N	Y	Y
Schoop U. <i>et al.</i> , 2007	N	N	Y	Y	N	Y	Y
Soukos NS. <i>et al.</i> , 2006	N	N	Y	Y	N	Y	Y
Vezzani MS. <i>et al.</i> , 2006	N	N	Y	Y	N	Y	Y
Bergmans L. <i>et al.</i> , 2006 <small>错误! 未定义书签。</small>	Y	Y	Y	Y	N	Y	Y
Perin FM. <i>et al.</i> , 2004	N	N	Y	Y	N	Y	Y
Kreisler M. <i>et al.</i> , 2003	N	N	Y	Y	N	Y	Y
Dostalova T. <i>et al.</i> , 2002	N	N	Y	Y	N	Y	Y
Schoop U. <i>et al.</i> , 2002	N	N	Y	Y	N	Y	Y
Piccolomini R. <i>et al.</i> , 2002 <small>错误! 未定义书签。</small>	Y	Y	Y	Y	N	Y	Y
Folwaczny M. <i>et al.</i> , 2002 <small>错误! 未定义书签。</small>	Y	Y	Y	Y	N	Y	Y
Moritz A. <i>et al.</i> , 1999	Y	Y	Y	Y	N	Y	Y
Mehl A. <i>et al.</i> , 1999	Y	Y	Y	Y	N	Y	Y

¹ Evidence-Based Medicine. How to Practise and Teach EBM, 2nd ed., Sackett D *et al.*, Churchill Livingstone, Edinburgh, UK.

² <http://www.cebm.net/index.aspx?o=1025>

³ Not applicable.

⁴ Also identified in Cochrane search.

Moritz A. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Ramskold LO. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Gutknecht N. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Moshonov J. <i>et al.</i> , 1995	N	N	Y	Y	N	Y	Y
Fegan SE. <i>et al.</i> , 1995	N	N	Y	Y	N	Y	Y
Hardee MW. <i>et al.</i> , 1994	N	N	Y	Y	N	Y	Y

2.3. Data Analysis

All results were converted and finally expressed in logCFU / ml (logarithm of the Colony Forming Units per milliliter) of the bacteria that were found after root canal preparation and laser or not laser irradiation. This was done for all specimens of all the studies.

Data from the studies that met validity criteria were extracted and classified by the method used for each bacterial group (Tables 15-21), for the polymicrobial studies (Table 22) and for all of the studies together (Table 23).

3. RESULTS

3.1. Medline Search

The MEDLINE search from 1966 to June 2010 identified 240 articles (Table 7). From 240 articles identified by the search, the hand examination of titles, abstracts and articles in full text revealed that 120 were irrelevant and 120 appeared to be relevant. Of the 120 relevant articles, 6 were *in vivo* studies, 5 expert's opinion, 6 narrative reviews, 3 case reports, 1 was an animal study and the remaining 99 were relevant *in vitro* studies.

Of the 99 relevant *in vitro* studies:

- Seventy were excluded because they were not related to the question.

Analytically

Seventeen examined the removal of smear layer and debris (错误! 未找到引用源。).

Twenty four examined morphological changes of the root canal (Table 9).

Twelve examined obturation and apical leakage after obturation (Table 10). Eleven examined dentin permeability (Table 11).

Five examined the thermal effects on the dentin (Table 12).

One examined adhesion of root canal sealers(5).

- One was excluded because it was not conducted on human teeth but on human teeth slices (6).
- Three were excluded because they did not provide quantitative results (7-9).

From the remaining 25 articles, 15 (10-24) were excluded because they met less than 5 of 7 validity criteria (错误! 未找到引用源。). Ten articles (25-34) were finally included. Details of the included 10 studies are presented in Table 14.

3.2. Cochrane Search

The Cochrane search identified 28 articles. From these 28 articles the hand examination of titles, abstracts and articles in full text, revealed that 16 were irrelevant to the question and 12 appeared to be relevant. Of the 12 arti-

cles, 7 did not meet the inclusion criteria and the remaining 5 were relevant *in vitro* studies. All of the 5 relevant *in vitro* studies were also identified by the Medline

Table 7. Search results.




Table 8. Excluded studies: Remove smear layer /debris.

Reference	Detail
Soares F. <i>et al.</i> , 2008	Remove smear layer
Moshonov J. <i>et al.</i> , 2004	Remove smear layer
Moshonov J. <i>et al.</i>, 2003	Remove smear layer
Matsuoka E. <i>et al.</i> , 1998	Remove smear layer
Takeda FH. <i>et al.</i> , 1998	Remove smear layer
Arrastia-Jitosho AM. <i>et al.</i> , 1998	Remove smear layer
Takeda FH. <i>et al.</i> , 1998	Remove smear layer
Radatti DA. <i>et al.</i> , 2006	Remove debris
Blum JY. <i>et al.</i> , 1997	Remove debris
Harashima T. <i>et al.</i> , 1997	Remove debris
Saunders WP. <i>et al.</i> , 1995	Remove debris
Machida T. <i>et al.</i> , 1995	Remove debris, Thermal effects
Moshonov J. <i>et al.</i> , 1995	Remove debris
Bahcall JK. <i>et al.</i> , 1993	Remove debris, Morphological

Frentzen M. <i>et al.</i> , 1991	Remove debris
Liesenhoff T. <i>et al.</i> , 1989	Remove debris, Morphological
Pini R. <i>et al.</i> , 1989	Remove debris

search, met all the inclusion criteria and 5 or more of the validity assessment criteria. So, they were finally included (**Table 6**).

3.3. Embase Search:

The Embase search from 1945 to June 2010 identified 35 articles. From these 35 articles identified by the search, the hand examination of titles, abstracts and articles in full text, revealed that all of them were irrelevant.

In order to be able to compare the bactericidal ability of the various treatment protocols that were applied on the human teeth *in vitro*, by each different group of researchers, the results were expressed in logCFU / ml. The CFU/ml of the bacteria that were found in the root canals after bacterial contamination and consecutive standard root canal preparation and / or laser irradiation was calculated and converted in logCFU / ml. The closer the

value of logCFU / ml is to 0, the greater the positive effect the treatment protocol has to the root canal disinfection.

LogCFU / ml of all bacteria is plotted in **Tables 15-23** (Clustered bars and 3-D clustered bars). On the X-axis lies the logCFU / ml and on the Y-axis the various treatment protocols.

Table 9. Excluded articles: Morphological changes.

Reference	Details
Gurbuz T. <i>et al.</i> , 2008	Morphological study
Da Costa Ribeiro A. <i>et al.</i> , 2007	Morphological, Thermal effects
Jahan KM. <i>et al.</i> , 2006	Morphological
Altundasar E. <i>et al.</i> , 2006	Morphological and Histochemical Changes
Matsuoka E. <i>et al.</i> , 2005	Morphological study
Biedma BM. <i>et al.</i> , 2005	Morphological study
Camargo SE. <i>et al.</i> , 2005	Morphological study
Niccoli-Filho W. <i>et al.</i> , 2005	Morphological study
Ali MN. <i>et al.</i> , 2005	Morphological study, Smear layer removal
Ishizaki NT. <i>et al.</i> , 2004	Morphological, Thermal effects
Khabbaz MG. <i>et al.</i> , 2004	Morphological study
Kesler G. <i>et al.</i> , 2002	Morphological study, Remove smear layer
Kaitsas V. <i>et al.</i> , 2001	Morphological and Histological changes
Matsuoka E. <i>et al.</i> , 2000	Morphological study
Yamazaki R. <i>et al.</i> , 2001	Morphological, Thermal effects
Barbakow F. <i>et al.</i> , 1999	Morphological study, Remove smear layer
Takeda FH. <i>et al.</i> , 1999	Morphological study, Remove smear layer
Eto JN. <i>et al.</i> , 1999	Morphological study, Remove smear layer
Takeda FH. <i>et al.</i> , 1998	Morphological study, Remove smear layer
Harashima T. <i>et al.</i> , 1998	Morphological study, Remove smear layer
Khan MA. <i>et al.</i> , 1997	Morphological, Thermal effects
Komori T. <i>et al.</i> , 1997	Morphological, Thermal effects
Goodis HE. <i>et al.</i> , 1993	Morphological study, Remove smear layer
Gutknecht N, Behrens VG, 1991	Morphological study

Table 10. Excluded articles: Obturation / Apical leakage.

Reference	Details
Ebihara A. <i>et al.</i> , 2002	Apical leakage after obturation
Kimura Y. <i>et al.</i> , 1999	Apical leakage after obturation
Kimura Y. <i>et al.</i> , 2001	Apical leakage after obturation
Park DS. <i>et al.</i> , 2001	Apical leakage after obturation
Goya C. <i>et al.</i> , 2000	Apical leakage after obturation, Removal of smear layer
Yamazaki R. <i>et al.</i> , 1999	Apical leakage after obturation

De Moura-Netto C. <i>et al.</i> , 2007	Apical sealing
Gekelman D. <i>et al.</i> , 2002	Apical sealing of root canal fillings
Carvalho CA. <i>et al.</i> , 2002	Apical sealing of root canal fillings.
Varella CH., Pillegi R., 2007	Obturation
Gharib SR. <i>et al.</i> , 2007	Obturation
Wang X. <i>et al.</i> , 2005	Obturation, Morphological, Thermal effects, Apical leakage

Table 11. Excluded articles: Dentin Permeability.

Reference	Details
Winik R. <i>et al.</i> , 2006	Marginal permeability, Tubular penetration,
Al-Azzawi LM. <i>et al.</i> , 2006	Dentin permeability
Aranha AC. <i>et al.</i> , 2005	Dentin permeability
Oliveira RG. <i>et al.</i> , 2004	Dentin permeability after apicoectomy
Gouw-Soares S. <i>et al.</i> , 2004	Dentin permeability
Arisu HD. <i>et al.</i> , 2004	Dentin permeability, Morphological
Brugnera A Jr. <i>et al.</i> , 2003	Dentin permeability
Lee BS. <i>et al.</i> , 2002	Dentin permeability
Pecora JD. <i>et al.</i> , 2000	Dentin permeability
Stabholz A. <i>et al.</i> , 1992	Dentin permeability
De Souza FD. <i>et al.</i> , 2005	Coronal microleakage

Table 12. Excluded articles: Thermal effects/ Adhesion of root canal sealers

Reference	Details
Nammour S. <i>et al.</i> , 2004	Thermal effects
Deutsch AS. <i>et al.</i> , 2004	Thermal effects
Amyra T. <i>et al.</i> , 2000	Thermal effects, Morphological
Cohen BI. <i>et al.</i> , 1996	Thermal effects
Neev J. <i>et al.</i> , 1993	Thermal effects

Table 13. Excluded Studies: not meeting validity criteria.

Reference	1	2	3	4	5	6	7
Fonseca MB. <i>et al.</i> , 2008	N	N	Y	Y	N	Y	Y
Gordon W. <i>et al.</i> , 2007	N	N	Y	Y	N	Y	Y
Schoop U. <i>et al.</i> , 2007	N	N	Y	Y	N	Y	Y
Soukos NS. <i>et al.</i> , 2006	N	N	Y	Y	N	Y	Y
Vezzani MS. <i>et al.</i> , 2006	N	N	Y	Y	N	Y	Y
Perin FM. <i>et al.</i> , 2004	N	N	Y	Y	N	Y	Y
Kreisler M. <i>et al.</i> , 2003	N	N	Y	Y	N	Y	Y
Dostalova T. <i>et al.</i> , 2002	N	N	Y	Y	N	Y	Y
Schoop U. <i>et al.</i> , 2002	N	N	Y	Y	N	Y	Y
Moritz A. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Ramskold LO. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Gutknecht N. <i>et al.</i> , 1997	N	N	Y	Y	N	Y	Y
Moshonov J. <i>et al.</i> , 1995	N	N	Y	Y	N	Y	Y
Fegan SE. <i>et al.</i> , 1995	N	N	Y	Y	N	Y	Y

Hardee MW. et al, 1994 N N Y Y N Y Y

• *Enterococcus faecalis*

Enterococcus faecalis survival was examined 19 times by 5 different groups of researchers.

A) Analysis of the results of studies with specific bacteria.

Table 14. Analysis of Treatment Protocols.

REFERENCES/ BACTERIA	STORAGE MATERIAL	IRRIGATION SOLUTION	INSTRUMENTATION	LASER	DURATION	NUMBER OF TEETH
de Souza EB et al (2008). <i>E. faecalis</i> (3*10 ⁸ cells/ml)	1% NaOCl	<u>G1,G2:</u> 0.5% NaOCl +urea peroxide cream+ 17% EDTA-T+ saline solution <u>G3:</u> Saline solution	G1, G2,: Rotary instrumentation (Crown down), No 50.	G1: diode laser(830nm/3W) G2: no laser irradiation G3: no laser irradiation (control)	2 days	<u>N = 30</u> G1 = 10 G2 = 10 G3 = 10
Fimpe JL et al (2008) <i>A.israelii</i> (2.5*10 ⁸ cells/ml) <i>F.nucleatum</i> (2.5*10 ⁸ cells/ml) <i>P.gingivalis</i> (2.5*10 ⁹ cells/ml) <i>P.intermedia</i> (2.5*10 ⁸ cells/ml)	0.5% NaOCl for 2 - 4 weeks.	<u>G1-G4:</u> RcPrep+ 6% NaOCl+ 17% EDTA+ 6%NaOCl	G1-G4: Rotary instrumentation (Crown-down) MAF: 0.465	<u>First set of experiments (n = 72.BHI broth):</u> G1: No laser/No MB G2: MB only G3: Diode laser (665 nm/ 1 W) G4: Diode laser (665 nm/1 W) and MB <u>Second set of experiments (n = 39.PBS broth):</u> G1: No laser/No MB G2: MB only G3: Diode laser (665 nm/ 1 W) and MB	7 days	<u>First set of experiments (n = 72):</u> G1 = 19 G2 = 18 G3 = 17 G4 = 18 <u>Second set of experiments (n = 39):</u> G1 = 13 G2 = 13 G3 = 13
Bergmans L. et al (2008) <i>S.anginosus</i> (4*10 ⁸ cells/ml)	0.5% chloramine in water at 4°C	<u>Ga-Gd:</u> 2.5% NaOCl + 17% EDTA + tap water	<u>Ga-Gd:</u> Rotary instrumentation (Crown down), No 30.	Ga: diode laser(635 nm/ 0.1 W)+ TBO+ RTF Gb: diode laser(635 nm/ 0.1 W) +RTF Gc: TBO +RTF Gd:RTF(positive control)	2 days	<u>N = 12</u> Ga = 3 Gb = 3 Gc = 3 Gd = 3
<i>E.faecalis:</i> (4*10 ⁸ cells/ml)	0.5% chloramine in water at 4°C	<u>Ga-Gd:</u> 2.5% NaOCl + 17% EDTA + tap water	<u>Ga-Gd:</u> Rotary instrumentation (Crown down), No 30.	Ga: diode laser(635 nm/ 0,1 W)+ TBO+ RTF Gb: diode laser(635 nm/ 0,1 W) +RTF Gc: TBO +RTF Gd:RTF(positive control)	2 days	<u>N = 12</u> Ga = 3 Gb = 3 Gc = 3 Gd = 3
<i>F.nucleatum</i> (4*10 ⁸ cells/ml)	0.5% chloramine in water at 4°C.	<u>Ga-Gd:</u> 2.5% NaOCl + 17% EDTA + tap water	<u>Ga-Gd:</u> Rotary instrumentation (Crown down), No 30.	Ga: diode laser(635 nm/ 0.1 W)+ TBO+ RTF Gb: diode laser(635 nm/ 0.1 W) +RTF Gc: TBO +RTF Gd:RTF(positive control)	2 days	<u>N = 12</u> Ga = 3 Gb = 3 Gc = 3 Gd = 3
Negative group (uninfected)	0.5% chloramine in water at 4°C	<u>Ga-Gd:</u> 2.5% NaOCl + 17% EDTA + tap water	<u>Ga-Gd:</u> Rotary instrumentation (Crown down), No 30.	No laser	2 days	N = 2
Foschi F, et al(2007) <i>E. faecalis</i> (5*10 ⁸ cells)	0.5% NaOCl for 2 weeks.	<u>G1-G4</u> -6% NaOCl: -17% EDTA deactivated with -6%NaOCl for 3 min.	<u>G1-G4</u> Rotary instrumentation (Crown down), MAF : 0.465	G1: no laser/no MB G2: MB G3: Diode laser (665 nm/1 W) G4: Diode laser (665 nm/1 W) and MB	6 days	<u>N = 64</u> G1 = 15 G2 = 15 G3 = 15 G4 = 15

Wang Q. <i>et al</i> (2007) E. faecalis ($> 10^8$ CFU/ml)	physiological saline solution at 4°C	G1-G4: 5.25% NaOCl+ physiological saline G5: 2.5% NaOCl + 5.25%NaHS+ physiological saline G6: None	G1-G6: Hand instrumentation (step-back), No 50	G1:Er,Cr:YSGG(2780 nm/ 1 W) G2:Er,Cr:YSGG(2780 nm/1,5W) G3:Nd:YAG(1064 nm/1W) G4:Nd:YAG(1064 nm/1,5W) G5:no laser irradiation (positive control) G6:no laser irradiation (negative control)	1 day	N = 60 G1 = 10 G2 = 10 G3 = 10 G4 = 10 G5 = 10 G6 = 10
Bergmans L. <i>et al</i> (2006) E. faecalis (4×10^8 CFU/ml)	0.5% chloramine in water at 4°C	G1,G2: 2.5% NaOCl+ 17% EDTA+ tap water + 0.9% sterile saline G3: 2.5% NaOCl +17% EDTA+ tap water	G1-G3: Rotary instrumentation (Crown down), No 30	G1: Nd:YAG laser (1064 nm/1.5 W/15 Hz)+RTF G2: RTF,no laser irradiation, Infected(positive control) G3: RTF, no laser irradiation, (uninfected, negative control)	2 days	N = 8 G1 = 3 G2 = 3 G3 = 2
Piccolomini R. <i>et al</i> (2002) A. naeslundii (1.8×10^8 CFU/ml)	none	SubA, SubB1, SubB2 Physiological saline + EDTA SubC: physiological saline+ EDTA+ 5.25% NaOCl	All groups: Hand instrumentation (Crown-down).	SubA: no laser irradiation (control) SubB1:Nd:YAG laser (1064 nm/5 Hz) Sub B2: Nd:YAG laser (1064 nm/ 10 Hz) SubC: no laser irradiation	1 day.	N = 60 SubA = 5 SubB1 = 10 SubB2 = 10 SubC = 5
Paeruginosa (1.8×10^8 CFU/ml)	none	SubA, SubB1, SubB2 Physiological saline + EDTA SubC: physiological saline+ EDTA+ 5.25% NaOCl	All groups: Hand instrumentation (Crown-down).	SubA: no laser irradiation (control) SubB1:Nd:YAG laser (1064 nm/ 5 Hz) Sub B2:Nd:YAG laser (1064 nm/10 Hz) SubC: no laser irradiation	1 day.	SubA = 5 SubB1 = 10 SubB2 = 10 SubC = 5
Folwaczny M. <i>et al</i> (2002) Escherichia Coli (8.67×10^6 CFU/ml)	Sterile saline solution	G1a-G1d: saline solution G1d: saline solution+ 1% NaOCl	G1a-G1d: Hand instrumentation, No 40.	G1a:No laser irradiation (positive control) G1b: Nd:YAG laser (1064 nm /0.005 W) G1c: Nd:YAG laser (1064 nm/ 0.01 W) G1d: No laser irradiation	1 day	N = 114 G1a = 13 G1b = 13 G1c = 13 G1d = 13
Folwaczny M. <i>et al</i> (2002) S. aureus (1.44×10^6 CFU/ml)	Sterile saline solution	G2a-G2d: saline solution G2d: saline solution+ 1% NaOCl	G2a-G2d: Hand instrumentation, No 40.	G2a:No laser irradiation(positive control) G2b: Nd:YAG laser (1064 nm /0.005 W) G2c: Nd:YAG laser (1064 nm/ 0.01 W) G2d: No laser irradiation	1 day	G2a = 13 G2b = 13 G2c = 13 G2d = 13
Negative control (uninfected)	Sterile saline solution	saline solution	Hand instrumentation, No 40.	G3: no laser irradiation (negative control)	1 day	G3 = 10

Moritz A, et al (1999) Escherichia Coli and Enterococcus faecalis (5*10 ⁵ CFU/mL)	Physiological saline solution	<u>All groups:</u> EDTA+ physiological saline solution	<u>All groups:</u> Rotary instrumentation, (Step-back)	G1: Positive control(untreated) G2: Nd:YAG laser (0.8 W/1064 nm) G3: Nd:YAG laser (1.5 W/1064 nm) G4: Ho: YAG laser (0.8 W/2130 nm) G5: Ho: YAG laser, (1.5 W/2130 nm) G6: Er:YAG laser, (0,8 W/2940 nm) G7: Er:YAG laser, (1.5 W/2940 nm)	1 day	N = 40 G1 : 10 G2 : 5 G3 : 5 G4 : 5 G5 : 5 G6 : 5 G7 : 5
Mehl A, et al (1999) Escherichia Coli (1.1*10 ⁶ CFU/ml)	none	<u>Ge1-Ge4:</u> sterile saline solution <u>Ge4:</u> 1.25% NaOCl	Hand instrumentation, No 40.	Gs1:no laser irradiation (positive control) Gs2:Er:YAG (2940 nm/0.003 W/15 sec/50 mJ) Gs3 : Er : YAG (2940 nm/0.001 W/60 sec/50 mJ) Gs4: no laser irradiation (1.25% NaOCl)	1 day	N = 90 Ge1 : 10 Ge2 : 10 Ge3 : 10 Ge4 : 10
S.aureus (1.53*10 ⁶ CFU/ml)	none	<u>Gs1-Gs4:</u> sterile saline solution <u>Gs4:</u> 1.25% NaOCl	Hand instrumentation, No 40.	Gs1:no laser irradiation (positive control) Gs2:Er:YAG (2940 nm/0.003 W/15 sec/50 mJ) Gs3 : Er : YAG (2940 nm/0.001 W/60 sec/50 mJ) Gs4: no laser irradiation (1.25% NaOCl)	1 day	Gs1 : 10 Gs2 : 10 Gs3 : 10 Gs4 : 10
Negative control (uninfected)	none	None	Hand instrumentation, No 40.	Gn: negative (uninfected)	1 day	Gn : 10

Table 15. Enterococcus faecalis survival.




Table 16. Streptococcus anginosus survival.





Table 17. Fusobacterium nucleatum survival.



In the first place, they contaminated the root canals with E.faecalis. In the second place they performed chemo-mechanical preparation and / or laser irradiation with or without specific solutions. The logCFU / ml ranges from 1,636 to 8,818 (Table 15).

Table 18. Escherichia coli survival.




Table 19. Actinomyces naeslundii survival.




Table 20. Staphylococcus aureus survival.





Table 21. Pseudomonas aeruginosa survival.



The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal and diode laser/ 665 nanometer / 1Watt irradiation with the combined effect

Table 22. Polymicrobial survival.





Table 23. Mean microbial survival.



of Methylene Blue (MB) as photosensitizing agent (logCFU / ml = 1,636) (33).

Instrumentation of the root canal and diode laser/ 665 nm / 1 W irradiation (logCFU / ml = 2,061) (33).

Instrumentation of the root canal combined with MB as photosensitizing agent (logCFU / ml = 2,190) (33).

Instrumentation followed by irrigation with 2, 5% NaOCl (sodium hypochloride)

(logCFU / ml = 2,602) (31).

When instrumentation and diode laser / 830 nm / 3 W were used, no CFU / ml was found. However, as the author explained, this result is due to the fact that the level of sensitivity of the methodology used, was insufficient for detecting viable cells in low concentrations (contact with the author via e-mail) (32).

- ***Streptococcus anginosus***

Streptococcus anginosus survival was examined 4 times by 1 group of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *S.anginosus*. Consequently, they completed root canal preparation and either used or not laser irradiation with specific solutions. The logCFU/ml ranges from 5,000 to 6,204 (**Table 16**) (30).

These results demonstrate no satisfying reduction of *S. anginosus* after diode laser / 635 nm / 0,1W irradiation combined with RTF (reduced transferred fluid) and / or TBO (toluidine blue), nor after instrumentation and irrigation with RTF and/or TBO (logCFU / ml > 5,000).

- ***Fusobacterium nucleatum***

Fusobacterium nucleatum survival was examined 4 times by 1 group of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *F.nucleatum*. Consequently, they completed root canal preparation and either used or not laser irradiation with specific solutions. The logCFU / ml ranges from 4,176 to 6,000 (**Table 17**) (30).

These results demonstrate no satisfying reduction of *F.nucleatum* after diode laser / 635 nm / 0,1 W irradiation combined with RTF (reduced transferred fluid) and / or TBO (toluidine blue), nor after instrumentation and irrigation with RTF and/or TBO (logCFU / ml > 4,176).

- ***Escherichia coli***

Escherichia coli survival was examined 8 times by 2 different groups of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *E. coli*. Consequently, they completed root canal preparation and either used or not laser irradiation with or without specific solutions. The logCFU/ml ranges from 2,342 to 6,938 (**Table 18**).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 1, 25% NaOCl

(logCFU / ml = 2,342) (28).

Instrumentation of the root canal and Er:YAG laser/ 2940 nm / 0,001 W irradiation

(logCFU / ml = 2,572) (28).

Instrumentation of the root canal followed by

irrigation with 1% NaOCl

(logCFU / ml = 3,012) (25).

Instrumentation of the root canal and Er:YAG laser/ 2940 nm / 0,003 W irradiation (logCFU / ml = 3,155) (28).

- ***Actinomyces naeslundii***

Actinomyces naeslundii survival was examined 4 times by 1 group of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *A. naeslundii*. Consequently, they completed root canal preparation and either used or not laser irradiation with or without specific solutions. The logCFU / ml ranges from 0 to 3,698 (

Table 19) (29).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 5, 25% NaOCl (logCFU / ml = 0).

Instrumentation of the root canal and Nd:YAG laser/ 1064 nm / 10 Hz / 15 sec irradiation (logCFU / ml = 3,053).

Instrumentation of the root canal and Nd:YAG laser/ 1064 nm / 5 Hz / 15sec irradiation (logCFU / ml = 3,518).

Instrumentation of the root canal solely (logCFU / ml = 3,698).

- ***Staphylococcus aureus***

Staphylococcus aureus survival was examined 8 times by 2 groups of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *S.aureus*. Consequently, they completed root canal preparation and either used or not laser irradiation with or without specific solutions. The logCFU / ml ranges from 2,703 to 6,184 (**Table 20**).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 1, 25% NaOCl

(logCFU / ml = 2,703) (28).

Instrumentation of the root canal and Er:YAG laser/ 2940 nm / 0,001 W irradiation (logCFU / ml = 2,973) (28).

Instrumentation of the root canal followed by irrigation with 1% NaOCl

(logCFU / ml = 3,246) (25).

Instrumentation of the root canal and Er:YAG laser/ 2940 nm / 0,003 W irradiation (logCFU / ml = 3,348) (28).

- ***Pseudomonas aeruginosa***

Pseudomonas aeruginosa survival was examined 4 times by 1 group of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with *P.aeruginosa*. Consequently, they completed root canal prepara-

tion and either used or not laser irradiation with or without specific solutions. The logCFU/ml ranges from 0 to 4,544 (**Table 21**) (29).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 5, 25% NaOCl (logCFU / ml = 0).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 10 Hz/ 15sec irradiation (logCFU / ml = 3,695).

B) Analysis of the Polymicrobial Studies Results

Microbial survival after polymicrobial infection of root canals was examined 30 times by 6 groups of researchers. In the first place they performed chemo-mechanical preparation and in the second place they contaminated the root canals with some of the following microorganisms; *E. coli*, *S. aureus*, *A. naeslundii*, *P. aeruginosa*, *S. anginosus*, *E. faecalis*, *F. nucleatum*, *A. israelii*, *P. gingivalis* and *P. intermedia*. Consequently, they completed root canal preparation and either used or not laser irradiation with or without specific solutions. The logCFU / ml ranges from 0 to 6,548 (**Table 22**).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 5,25 %NaOCl (logCFU / ml = 0) (29).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,8 W irradiation

(logCFU / ml = 1,924) (27).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 1,5W irradiation

(logCFU / ml = 2,113) (27).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 1,5W irradiation

(logCFU / ml = 2,477) (27).

Instrumentation of the root canal and Ho: YAG laser/ 2130 nm / 0,8 W irradiation

(logCFU / ml = 2,491) (27).

Instrumentation of the root canal followed by irrigation with 1,25% NaOCl (logCFU / ml = 2,522) (28).

Instrumentation of the root canal and Ho: YAG laser/ 2130 nm / 1,5W irradiation

(logCFU / ml = 2,531) (27).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,001W irradiation

(logCFU / ml = 2,772) (28).

Instrumentation of the root canal followed by irrigation with 1% NaOCl (logCFU / ml = 3,138) (25).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 0,8 W irradiation

(logCFU / ml = 3,204) (27).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,003W irradiation

(logCFU / ml = 3,251) (28).

Instrumentation of the root canal and diode laser/ 665 nm / 1W irradiation combined with MB as photosensitizing agent and PBS (Phosphate-Buffered Saline) broth (logCFU / ml = 3,341) (34).

Instrumentation of the root canal and diode laser/ 665 nm / 1W irradiation combined with MB as photosensitizing agent (logCFU / ml = 3,344) (34).

Instrumentation of the root canal and Nd: YAG laser / 1064 nm / 10 Hz / 15sec irradiation(logCFU / ml = 3,374) (29).

Instrumentation of the root canal combined with MB as photosensitizing agent (logCFU / ml=3,814) (34).

Instrumentation of the root canal combined with MB as photosensitizing agent and PBS (Phosphate-Buffered Saline) broth (logCFU / ml = 3,906) (34).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 5 Hz / 15sec irradiation (logCFU / ml = 3,993) (29).

C) Analysis of the Mean Averages of Microbial Survival in general

The survival of 10 microorganisms well connected with endodontic problems: *E. faecalis*, *S. anginosus*, *F. nucleatum*, *A. israelii*, *P. gingivalis*, *P. intermedia*, *A. naeslundii*, *P. aeruginosa*, *E. coli*, and *S. aureus* was examined by 10 groups of researchers regarding the bactericidal effect of 30 different treatment protocols on them. In the first place the majority of them performed chemo-mechanical preparation and in the second place they contaminated the root canals with one or more microorganisms. Then, they completed the root canal preparation and either used or not laser irradiation with or without specific solutions. The logCFU / ml ranges from 0 to 7,940 (**Table 23**).

The best treatment protocols for root canal disinfection in descending order are as follows:

Instrumentation of the root canal followed by irrigation with 5,25% NaOCl (logCFU / ml = 0) (29).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,8 W irradiation

(logCFU / ml = 1,924) (27).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 1,5 W irradiation

(logCFU / ml = 2,133) (27).

Instrumentation of the root canal and diode laser/ 665 nm / 1W irradiation with the combined effect of Methylene Blue (MB) as photosensitizing agent (logCFU / ml = 2,490) (33, 34).

Instrumentation of the root canal and Ho: YAG laser/ 2130 nm / 0,8W irradiation

(logCFU / ml = 2,491) (27).

Instrumentation of the root canal followed by irrigation with 1,25% NaOCl (logCFU / ml = 2,522) (28).

Instrumentation of the root canal and Ho: YAG laser/ 2130 nm / 1,5W irradiation (logCFU / ml = 2,531) (27).

Instrumentation followed by irrigation with 2, 5% NaOCl (logCFU / ml = 2,602) (31).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,001W irradiation (logCFU / ml = 2,772) (28).

Instrumentation of the root canal with the combined effect of Methylene Blue (MB) as photosensitizing agent (logCFU / ml = 3,002) (33, 34).

Instrumentation of the root canal and diode laser/ 665 nm / 1W irradiation (logCFU / ml = 3,033) (33, 34).

Instrumentation followed by irrigation with 1% NaOCl (logCFU / ml = 3,138) (25).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 0, 8 W irradiation (logCFU / ml = 3,204) (27).

Instrumentation of the root canal and Er: YAG laser/ 2940 nm / 0,003 W irradiation (logCFU / ml = 3,251) (28).

Instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 10 Hz / 15 sec irradiation (logCFU / ml = 3,374) (29).

Instrumentation of the root canal and Nd: YAG laser / 1064 nm / 5 Hz/ 15 sec irradiation (logCFU / ml = 3,993) (29).

4. DISCUSSION

Generally, it is well known that certain microorganisms are related to specific pathological situations in endodontics. Being aware of them is necessary so as to be able to consider the importance of the disinfection capacity of each treatment plan. Consequently, it has been proved that:

- 1) With symptomatic endodontic disease and apical bone resorption *T. denticola* is associated (1).
- 2) In endodontically infected teeth without a sinus tract *E. faecalis* and *Strept. anginosus* were mostly found (2).
- 3) In teeth with necrotic pulp *P. gingivalis*, *P. endodontalis*, *P. intermedia*, and *P. nigrescens* were identified more frequently(4).
- 4) With root canal treatment failures *E. faecalis* is associated (1,3).
- 5) In endodontically infected teeth with sinus tracts *P. gingivalis* and *F. nucleatum* were mostly identified (2).

This systematic review identified 10 *in vitro* studies that examined the effectiveness of treatment protocols, among classical methods and/or various laser applications in root canal disinfection. The results of these 10

studies indicated that the treatments which provide the best bactericidal ability regarding each bacterial solely and all of them were, in descending order of efficacy:

4.1. Enterococcus Faecalis

Instrumentation of the root canal and diode laser / 665 nm / 1 W irradiation with the combined effect of Methylene Blue (MB) as photosensitizing agent seemed to be better in root canal disinfection than instrumentation of the root canal and diode laser / 665 nm / 1 W irradiation. In addition, the last method was slightly better than instrumentation of the root canal combined with MB as photosensitizing agent and significantly better than instrumentation followed by irrigation with 2, 5% NaOCl (sodium hypochloride).

Regarding the instrumentation and diode laser / 830 nm / 3 W irradiation where no CFU / ml was found, this is due to the lack of sensitivity of the methodology used to detect low concentrations of viable cells (contact with author).

4.2. Streptococcus Anginosus

Instrumentation of the root canal and diode laser/ 635 nm / 0,1 W irradiation combined with RTF (reduced transferred fluid) and / or TBO (toluidine blue), as well as instrumentation with RTF and /or TBO demonstrated neither accepted nor satisfying reduction of *S.anginosus* (logCFU / ml = 5,000, logCFU / ml = 6,079, logCFU / ml = 6,113 and logCFU / ml = 6,204, respectively).

4.3. Fusobacterium Nucleatum

Instrumentation of the root canal and diode laser/ 635 nm / 0,1W irradiation combined with RTF (reduced transferred fluid) and/or TBO (toluidine blue), as well as instrumentation with RTF and /or TBO showed insufficient disinfection of *F.nucleatum*. (logCFU / ml = 4,176, log- CFU / ml = 6, logCFU / ml = 5,939and logCFU / ml = 5,986, respectively).

4.4. Escherichia Coli

Instrumentation of the root canal followed by irrigation with 1, 25% NaOCl seemed to be better in root canal disinfection than instrumentation of the root canal and Er:YAG laser/2940nm/0,001W irradiation. Furthermore, instrumentation of the root canal followed by irrigation with 1%NaOCl was less effective than the previous treatments, but slightly better than instrumentation of the root canal and Er: YAG laser/2940nm/0,003W irradiation.

4.5. Actinomyces Naeslundii:

Instrumentation of the root canal followed by irrigation with 5, 25% NaOCl seems to be the best in root canal disinfection as with this concentration no viable cells of

Actinomyces naeslundii are detected. This method is used and mentioned only by one group of researchers (29). Concerning the other treatments, they are significantly worse and in fact instrumentation of the root canal and Nd: YAG laser / 1064 nm / 10 Hz / 15sec irradiation is slightly better than the same method but with 5 Hz irradiation or instrumentation solely.

4.6. Staphylococcus Aureus

Instrumentation of the root canal followed by irrigation with 1, 25% NaOCl was better enough than instrumentation and Er: YAG laser / 2940 nm / 0,001W irradiation and seemed to be better in root canal disinfection than instrumentation of the root canal followed by irrigation with 1%NaOCl and instrumentation with Er: YAG laser / 2940nm/0,003W irradiation.

4.7. Pseudomonas aeruginosa:

Instrumentation of the root canal followed by irrigation with 5, 25% NaOCl was the best in root canal disinfection, as no CFU / ml of Pseudomonas aeruginosa was found. This method was conducted by one group of researchers (29). Regarding instrumentation and Nd: YAG laser / 1064 nm / 10 Hz / 15sec irradiation, it was moderately worse than the previous one.

4.8. Crobial Studies Results:

Instrumentation of the root canal followed by irrigation with 5, 25% NaOCl seemed to be the best in root canal disinfection as no viable cells were detected. However, this concentration of NaOCl is not used *in vivo* because it actively attacks living tissue without contributing significantly to treatment (35). Furthermore, instrumentation and Er: YAG laser/2940nm/0,8W irradiation was well successful and slightly better than the same treatment but with 1,5Watt irradiation, which was also better than instrumentation of the root canal and Nd: YAG laser/ 1064 nm / 1,5W irradiation. Also, instrumentation of the root canal and Ho: YAG laser/2130nm/0,8W irradiation was effective enough, as well as instrumentation of the root canal followed by irrigation with 1, 25%NaOCl and instrumentation followed with Ho: YAG laser/ 2130 nm / 1,5 W irradiation. Finally, instrumentation of the root canal and Er: YAG laser / 2940 nm / 0,001W irradiation was less effective than the previous methods. All the other methods seemed to be worse in root canal disinfection.

4.7. Microbial Survival in general:

Instrumentation of the root canal followed by irrigation with 5, 25%NaOCl seemed to be the best in root canal disinfection as no viable cells were detected. Instrumentation and Er: YAG laser / 2940 nm / 0,8W

irradiation was well successful and slightly better than the same treatment but with 1,5Watt irradiation. Instrumentation followed with diode laser / 665 nm / 1W irradiation with the combined effect of Methylene Blue (MB) as photosensitizing agent showed the same results when compared to instrumentation of the root canal and Ho: YAG laser / 2130 nm / 0,8W irradiation. Also good results show instrumentation of the root canal followed by irrigation with 1,25%NaOCl and Ho: YAG laser/ 2130 nm / 1,5 W irradiation which both are slightly better than instrumentation followed by irrigation with 2, 5% NaOCl and Er: YAG laser / 2940 nm / 0,001W irradiation. All the other methods seemed to be worse in root canal disinfection.

5. CONCLUSION

There are treatment protocols with the assistance or not of laser irradiation that can eliminate E.faecalis, E.coli and S.aureus inside the root canal. However, there is a serious number of S.anginosus, F.nucleatum, A. naeslundii and P.aeruginosa that remain inside the root canal even after laser irradiation. *In vitro*, NaOCl 5% seems to be the strongest solution in root canal disinfection. Concluding, it seems that new research is needed in order to set a treatment protocol effective in the root canal disinfection from all bacteria mentioned above that are related to endodontic origin pathology.

REFERENCES

- [1] Foschi F, Cavrini F, Montebugnoli L, Stashenko P, Sambri V, Prati C. Detection of bacteria in endodontic samples by polymerase chain reaction assays and association with defined clinical signs in Italian patients. *Oral Microbiol Immunol.* 2005 20(5):289-295.
- [2] Sassone L, Fidel R, Faveri M, Fidel S, Figueiredo L, Feres M. Microbiological evaluation of primary endodontic infections in teeth with and without sinus tract. *Int Endod J.* 2008;41(6):508-515.
- [3] Gomes B, Pinheiro E, Gadê-Neto C, Sousa E, Ferraz C, Zaia A, et al. Microbiological examination of infected dental root canals. *Oral Microbiol Immunol.* 2004;19(2):71-76.
- [4] Gomes B, Jacinto R, Pinheiro E, Sousa E, Zaia A, Ferraz C, et al. Porphyromonas gingivalis, Porphyromonas endodontalis, Prevotella intermedia and Prevotella nigrescens in endodontic lesions detected by culture and by PCR. *Oral Microbiol Immunol.* 2005;20(4):211-215.
- [5] Sousa-Neto M, Marchesan M, Pécora J, Junior A, Silva-Sousa Y, Saquy P. Effect of Er:YAG laser on adhesion of root canal sealers. *Journal of Endodontics* 2002 28(3):185-187.
- [6] Schoop U, Kluger W, Dervisebegovic S, Goharkhay K, Wernisch J, Georgopoulos A, et al. Innovative Wavelengths in Endodontic Treatment. *Lasers Surg. Med.* 2006;38(6):624-630.
- [7] Jha D, Guerrero A, Ngo T, Helfer A, Hasselgren G.

- Inability of laser and rotary instrumentation to eliminate root canal infection. *J Am Dent Assoc.* 2006 137(1):67-70.
- [8] Garcez A, Ribeiro M, Tegos G, Núñez S, Jorge A, Hamblin M. Antimicrobial photodynamic therapy combined with conventional endodontic treatment to eliminate root canal biofilm infection. *Lasers Surg Med* 2007;39(1):59-66.
- [9] Blum J, Michailesco P, Abadie M. An evaluation of the bactericidal effect of the Nd:YAP laser. *Journal of Endodontics* 1997 23(9):583-585.
- [10] 10. Dostálová T, Jelínková H, Housová D, Sulc J, Nemeč M, Dusková J, et al. Endodontic treatment with application of Er:YAG laser waveguide radiation disinfection. *J Clin Laser Med Surg.* 2002 20(3):135-139.
- [11] Fegan S, Steiman H. Comparative evaluation of the antibacterial effects of intracanal Nd:YAG laser irradiation: an in vitro study. *Journal of Endodontics* 1995 21(8):415-417.
- [12] Fonseca M, Júnior P, Pallota R, Filho H, Denardin O, Rapoport A, et al. Photodynamic therapy for root canals infected with *Enterococcus faecalis*. *Photomed Laser Surg.* 2008 26(3):209-213.
- [13] Gordon W, Atabakhsh V, Meza F, Doms A, Nissan R, Rizoio I, et al. The antimicrobial efficacy of the erbium, chromium:yttrium-scandium-gallium-garnet laser with radial emitting tips on root canal dentin walls infected with *Enterococcus faecalis*. *J Am Dent Assoc.* 2007;138(7):992-1002.
- [14] Gutknecht N, Nuebler-Moritz M, Burghardt S, Lampert F. The efficiency of root canal disinfection using a holmium:yttrium-aluminum-garnet laser in vitro. *J Clin Laser Med Surg.* 1997;15(2):75-78.
- [15] Hardee M, Miserendino L, Kos W, Walia H. Evaluation of the antibacterial effects of intracanal Nd:YAG laser irradiation. *Journal of Endodontics* 1994;20(8):377-380.
- [16] Kreisler M, Kohlen W, Beck M, Al Haj H, Christoffers A, Götz H, et al. Efficacy of NaOCl/H₂O₂ irrigation and GaAlAs laser in decontamination of root canals in vitro. *Lasers Surg Med.* 2003;32(3):189-196.
- [17] Moritz A, Gutknecht N, Goharkhay K, Schoop U, Wernisch J, Sperr W. In vitro irradiation of infected root canals with a diode laser: results of microbiologic, infrared spectrometric, and stain penetration examinations. *Quintessence Int.* 1997 28(3):205-209.
- [18] Moshonov J, Orstavik D, Yamauchi S, Pettiette M, Trope M. Nd:YAG laser irradiation in root canal disinfection. *Endod Dent Traumatol.* 1995 11(5):220-224.
- [19] Perin F, França S, Silva-Sousa Y, Alfredo E, Saquy P, Estrela C, et al. Evaluation of the antimicrobial effect of Er:YAG laser irradiation versus 1% sodium hypochlorite irrigation for root canal disinfection. *Aust Endod J.* 2004;30(1):20-22.
- [20] Ramsköld L, Fong C, Strömberg T. Thermal effects and antibacterial properties of energy levels required to sterilize stained root canals with an Nd:YAG laser. *Journal of Endodontics* 1997 23(2):96-100.
- [21] Schoop U, Goharkhay K, Klimscha J, Zagler M, Wernisch J, Georgopoulos A, et al. The use of the erbium, chromium:yttrium-scandium-gallium-garnet laser in endodontic treatment: the results of an in vitro study. *J Am Dent Assoc.* 2007;138(7):949-955.
- [22] Schoop U, Moritz A, Kluger W, Patruta S, Goharkhay K, Sperr W, et al. The Er:YAG laser in endodontics: results of an in vitro study. *Lasers Surg Med.* 2002;30(5):360-364.
- [23] Soukos N, Chen P, Morris J, Ruggiero K, Abernethy A, Som S, et al. Photodynamic therapy for endodontic disinfection. *Journal of Endodontics* 2006 32(10):979-984.
- [24] Vezzani M, Pietro R, Silva-Sousa Y, Brugnera-Junior A, Sousa-Neto M. Disinfection of root canals using Er:YAG laser at different frequencies. *Photomed Laser Surg.* 2006;24(4):499-502.
- [25] Folwaczny M, Mehl A, Jordan C, Hickel R. Antibacterial Effects of Pulsed Nd:YAG Laser Radiation at Different Energy Settings in Root Canals. *Journal of Endodontics* 2002;28(1):24-29.
- [26] Bergmans L, Moisiadis P, Teughels W, Van Meerbeek B, Quirynen M, Lambrechts P. Bactericidal effect of Nd:YAG laser irradiation on some endodontic pathogens ex vivo. *International Endodontic Journal* 2006;39(7):547-557.
- [27] Moritz A, Schoop U, Goharkhay K, Jakolitsch S, Kluger W, Wernisch J, et al. The bactericidal effect of Nd:YAG, Ho:YAG, and Er:YAG laser irradiation in the root canal: an in vitro comparison. *J Clin Laser Med Surg* 1999;17(4):161-164.
- [28] Mehl A, Folwaczny M, Haffner C, Hickel R. Bactericidal Effects of 2.94 μm Er:YAG-Laser Radiation in Dental Root Canals. *Journal of Endodontics* 1999;25(7):490-493.
- [29] Piccolomini R, D'Arcangelo C, D'Ercole S, Catamo G, Schiaffino G, De Fazio P. Bacterologic Evaluation of the Effect of Nd:YAG Laser Irradiation in Experimental Infected Root Canals. *Journal of Endodontics* 2002;28(4):276-278.
- [30] Bergmans L, Moisiadis P, Huybrechts B, Van Meerbeek B, Quirynen M, P. L. Effect of photo-activated disinfection on endodontic pathogens ex vivo. *International Endodontic Journal* 2008;41(3):227-239.
- [31] Wang Q, Zhang C, Yin X. Evaluation of the bactericidal effect of Er,Cr:YSGG, and Nd:YAG lasers in experimentally infected root canals. *Journal of Endodontics* 2007;33(7):830-832.
- [32] de Souza E, Cai S, Simionato M, Lage-Marques J. High-power diode laser in the disinfection in depth of the root canal dentin. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008 106(1):68-72.
- [33] Foschi F, Fontana C, Ruggiero K, Riahi R, Vera A, Doukas A, et al. Photodynamic Inactivation of *Enterococcus faecalis* in Dental Root Canals In Vitro. *Lasers Surg. Med.* 2007;39(10):782-787.
- [34] Fimple J, Fontana C, Foschi F, Ruggiero K, Song X, Pagonis T, et al. Photodynamic Treatment of Endodontic Polymicrobial Infection In Vitro. *Journal of Endodontics* 2008;34(6):728-734.