

# Journal of Dental Research

<http://jdr.sagepub.com>

---

## **Diversity of Endodontic Microbiota Revisited**

J.F. Siqueira, Jr. and I.N. Rôças  
*J DENT RES* 2009; 88; 969  
DOI: 10.1177/0022034509346549

The online version of this article can be found at:  
<http://jdr.sagepub.com/cgi/content/abstract/88/11/969>

---

Published by:



<http://www.sagepublications.com>

On behalf of:

International and American Associations for Dental Research

**Additional services and information for *Journal of Dental Research* can be found at:**

**Email Alerts:** <http://jdr.sagepub.com/cgi/alerts>

**Subscriptions:** <http://jdr.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

J.F. Siqueira Jr.\* and I.N. Rôças

Department of Endodontics and Molecular Microbiology, Faculty of Dentistry, Estácio de Sá University, Av. Alfredo Baltazar da Silveira, 580/cobertura, Recreio, Rio de Janeiro, RJ, Brazil 22790-710; \*corresponding author, [jf\\_siqueira@yahoo.com](mailto:jf_siqueira@yahoo.com) or [siqueira@estacio.br](mailto:siqueira@estacio.br)

*J Dent Res* 88(11):969–981, 2009

## ABSTRACT

Although fungi, archaea, and viruses contribute to the microbial diversity in endodontic infections, bacteria are the most common micro-organisms occurring in these infections. Datasets from culture and molecular studies, integrated here for the first time, showed that over 460 unique bacterial taxa belonging to 100 genera and 9 phyla have been identified in different types of endodontic infections. The phyla with the highest species richness were *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, and *Proteobacteria*. Diversity varies significantly according to the type of infection. Overall, more taxa have been disclosed by molecular studies than by culture. Many cultivable and as-yet-uncultivated phylotypes have emerged as candidate pathogens based on detection in several studies and/or high prevalence. Now that a comprehensive inventory of the endodontic microbial taxa has been established, future research should focus on the association with different disease conditions, functional roles in the community, and susceptibility to antimicrobial treatment procedures.

**KEY WORDS:** endodontic microbiology, molecular biology methods, culture, taxonomy.

DOI: 10.1177/0022034509346549

Received March 28, 2008; Last revision October 1, 2008; Accepted November 4, 2008; Updated July 3, 2009

A supplemental appendix to this article is published electronically only at <http://jdr.sagepub.com/supplemental>.

# Diversity of Endodontic Microbiota Revisited

## INTRODUCTION

Essentially, endodontic infection is the infection of the dental root canal system and the major etiologic agent of apical periodontitis (Siqueira, 2008). Although chemical and physical factors can induce periradicular inflammation, a large body of scientific evidence clearly indicates that micro-organisms are essential to the progression and perpetuation of the different forms of apical periodontitis (Kakehashi *et al.*, 1965; Sundqvist, 1976; Möller *et al.*, 1981). For endodontic infection to develop, the root canal must be devoid of vital pulp tissue and its defenses, as a consequence of either pulp necrosis (as a sequel to caries, trauma, periodontal disease, or iatrogenic operative procedures) or pulp removal for treatment. The borderline between the infecting microbiota and the host defenses is often located intraradicularly, *i.e.*, short of or at the apical foramen. In some cases, however, micro-organisms may reach the periradicular tissues, and the borderline is then situated extraradicularly, *i.e.*, beyond the boundaries of the apical foramen.

Because apical periodontitis is an infectious disease, the rationale for endodontic treatment is inarguably to eradicate the occurring infection and/or to prevent micro-organisms from infecting or re-infecting the root canal or the periradicular tissues. The cardinal principle of any healthcare profession is the thorough understanding of the disease etiology and pathogenesis, which provides a framework for effective prevention and treatment. In this context, understanding and defining the endodontic microbiota associated with different forms of disease are the basis for an endodontic practice of high quality and founded on a solid scientific basis. Such knowledge has the potential to contribute to the development of more effective preventive and therapeutic protocols. Furthermore, establishing an inventory of endodontic bacteria may also help future investigations of potential sources of pathogenic species for diseases in other human sites.

Traditionally, endodontic infections have been studied by means of culture approaches. Such studies have resulted in the establishment of a set of species thought to play an important role in the pathogenesis of apical periodontitis. More recently, not only have findings from culture-based methods been confirmed, but they have also been significantly supplemented with those from culture-independent molecular diagnostic techniques. Molecular methods have confirmed and strengthened the association of many cultivable bacterial species with apical periodontitis and have also revealed suspected new endodontic pathogens (Siqueira and Rôças, 2005b). The list of candidate pathogens has expanded to include culture-difficult species or even as-yet-uncultivated bacteria that had never been previously found in endodontic infections by culturing approaches. As a consequence, the endodontic microbiota has been refined and redefined by molecular methods (Siqueira and Rôças, 2005b).

Because knowledge of the endodontic infection associated with different clinical conditions originates from culture-dependent and culture-independent molecular biology studies, the most comprehensive view of the diversity of the endodontic microbiota requires integration of the data from both types of studies. This review focuses on the diversity of the microbiota in different types of endodontic infections and is mostly based on cataloguing the microbial taxa already identified in culture and molecular studies. Overall, data from 128 studies published up to March 2008, were used for compilation (listed in the Appendix). Culture studies included in this compilation are restricted to those performed from the mid-1970s to 2008, when reliable methods for the cultivation and identification of fastidious anaerobic bacteria were introduced and adopted in endodontic microbiology research. Species names were considered valid when listed on the DSMZ Bacterial Nomenclature Web site ([http://www.dsmz.de/microorganisms/bacterial\\_nomenclature.php](http://www.dsmz.de/microorganisms/bacterial_nomenclature.php)) or mentioned in recent articles published in the *International Journal of Systematic and Evolutionary Microbiology*.

## MOLECULAR STUDIES AND THE UNCULTIVATED MAJORITY

Culture methods have been highly successful over the past 100 years, nurturing the expectation that most micro-organisms can be grown in the laboratory. It is now widely recognized that culture techniques can strongly bias and underestimate the diversity of microbial populations (Hugenholtz, 2002; Relman, 2002). Only a small fraction of the bacteria present in most microbial ecosystems is amenable to propagation *ex vivo* (Handelsman, 2004; Fredricks and Marrazzo, 2005). Less than 1% of bacteria have been estimated to be cultivated from environments such as sea water, lakes, and soil (Amann *et al.*, 1995; Gewin, 2006). To sidestep the limitations of culture, tools and procedures based on molecular biology have become available and have been substantially improved to achieve a more realistic description of the microbial world without the need for cultivation. The most significant contributions of culture-independent molecular biology methods to medical microbiology relate to the identification of previously unknown human pathogens (Fredricks and Relman, 1996; Relman, 1999; Kellam and Weiss, 2001; Lawson, 2004) and the discovery of a far broader diversity of the human microbiota associated with different human sites. Studies based on the 16S rRNA gene approach have re-vealed that about 60% of the bacterial species in the oral cavity (Paster *et al.*, 2001; Aas *et al.*, 2005; Kumar *et al.*, 2005; de Lillo *et al.*, 2006), 50% on the skin (Dekio *et al.*, 2005), 38% in the esophagus (Pei *et al.*, 2004), 50% in the stomach (Bik *et al.*, 2006), 45% in the vagina (Fredricks *et al.*, 2005), and about 80% in the gut (Suau *et al.*, 1999; Eckburg *et al.*, 2005) represent as-yet-uncultivated and uncharacterized bacteria.

The study of the human gastrointestinal microbiota is a formidable example of the revolution fomented by molecular studies with regard to the study of the human microbiota in health and disease (Furrie, 2006). Culture-dependent studies of the gastrointestinal microbiota have resulted in the

characterization of over 400 species (Rajilić-Stojanović *et al.*, 2007). Application of molecular biology techniques revealed that the microbiota in the gut is significantly more complex than previously anticipated, since only a fraction of the bacteria encountered can currently be cultivated (Suau *et al.*, 1999; Eckburg *et al.*, 2005). Another major outcome of the revolution brought about by molecular methods was the recognition that the composition of the microbiota is subject-specific and dominated by as-yet-uncultivated and uncharacterized phylotypes (Hold *et al.*, 2002; Eckburg *et al.*, 2005; Hayashi *et al.*, 2005). The integration of data from culture and molecular studies reveals that more than 1000 different taxa compose the microbial diversity in the human gastrointestinal tract (Rajilić-Stojanović *et al.*, 2007).

The oral cavity is another good example of the impact of culture-independent molecular biology methods on the knowledge of the microbial diversity. Data from culture and molecular studies have collectively revealed that almost 800 distinct bacterial taxa may be able to live in the human oral cavity, though not all of them are present in the same individual at the same time (Paster *et al.*, 2006). Indeed, any particular individual can harbor about 100-200 of these 800 taxa of oral bacteria, indicating that there is a substantial diversity among different people (Paster *et al.*, 2006). Taken as a whole, bacteria detected from the oral cavity fall into 13 separate phyla, namely, *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, *Spirochaetes*, *Fusobacteria*, *Synergistes* (including phylotypes previously assigned to the phylum *Deferribacteres*), SR1 (originating from the OP11 division), TM7, *Chloroflexi*, *Deinococcus*, *Acidobacteria*, and *Cyanobacteria* (Paster *et al.*, 2001, 2002; Kazor *et al.*, 2003; de Lillo *et al.*, 2006; Aas *et al.*, 2007). This number may be even higher, since a recent study with DNA microarray technology suggested that members of 4 other phyla (*Aquificae*, *Nitrospira*, *Planctomycetes*, and *Thermomicrobia*) may have oral representatives, even though none has been identified (Huyghe *et al.*, 2008). A recent study with pyrosequencing, a high-throughput molecular approach that allows for extensive sequencing of microbial populations, revealed about 5600 and 10,000 species-level phylotypes representing 22 phyla in saliva and plaque, respectively (Keijsers *et al.*, 2008). The estimated number of oral phylotypes is about 20,000, which is considerably higher when compared with that reported in previous culture and clone library studies.

It is worth pointing out that both culture and molecular approaches have their own limitations in portrayals of microbial diversity. Limitations of culture have been extensively recognized, and many of them have been successfully overcome by molecular technology (Fredricks and Relman, 1999; Hugenholtz, 2002; Siqueira and Rôças, 2005a). However, molecular methods can also give a biased picture of bacterial diversity, and their limitations in this regard are mostly related to different levels of effectiveness of cell lysis for DNA extraction and PCR-related biases (von Wintzingerode *et al.*, 1997; Siqueira and Rôças, 2005a; Rajilić-Stojanović *et al.*, 2007).

It has been claimed that the high sensitivity of molecular methods allows for the detection of species present in very low

numbers and, consequently, with possibly no clinical importance. However, in microbial ecology, it has been demonstrated that some of these low-abundant bacteria might still serve as keystone species within complex mixed consortia (Sogin *et al.*, 2006). Moreover, some of the low-abundant species may be a result of historical ecological changes, in the sense that they may have been dominant in the past or have the potential to become dominant in the future in response to shifts in environmental conditions that favor their growth. Thus, low-abundant populations may eventually become dominant in response to environmental changes (Sogin *et al.*, 2006). Furthermore, species that are low-abundant in one individual can dominate the community in another individual. Therefore, at least from an ecological perspective, all species in a mixed community should be successfully detected and identified (Siqueira and Rôças, 2009b).

A broad range of molecular biology techniques has been used for diverse purposes in microbial ecology research (Siqueira and Rôças, 2005a; Spiegelman *et al.*, 2005). In endodontic microbiology research, the vast majority of molecular studies have addressed the issue of species composition in different types of endodontic infections. In spite of the great advantages of these methods, at this time, there is no available technique that can provide a comprehensive list of all bacterial species in a sample. As a consequence, a variety of techniques has been used to offer a better picture of the endodontic bacterial communities.

The chronology of the study of the diversity of the endodontic microbiology can be didactly divided into phases based on different strategic approaches. Early studies of the endodontic microbiota were conducted with broad-range culture methods (Bergenholtz, 1974; Kantz and Henry, 1974; Wittgow and Sabiston, 1975; Sundqvist, 1976; Baumgartner and Falkler, 1991; Sundqvist, 1992). These were followed by a generation of studies that used molecular detection methods, such as species-specific PCR and the original checkerboard DNA-DNA hybridization assay, to target cultivable bacteria previously isolated from infected canals or from other diseased oral sites (Conrads *et al.*, 1997; Jung *et al.*, 2000; Machado de Oliveira *et al.*, 2000; Rupf *et al.*, 2000; Siqueira *et al.*, 2000a,b, 2001b). These methods allowed for the inclusion of some culture-difficult species in the set of putative endodontic pathogens. The adoption of 16S rRNA gene clone library analysis facilitated an even more comprehensive broad-range investigation of bacterial communities in endodontic infections (Munson *et al.*, 2002; Saito *et al.*, 2006; Sakamoto *et al.*, 2006, 2008). By this approach, not only cultivable species but also as-yet-uncultivated and uncharacterized bacteria have been identified. Studies with 16S rRNA gene clone library analysis have revealed that 40-55% of the bacterial taxa found in primary endodontic infections have not been cultivated and validly named (Munson *et al.*, 2002; Sakamoto *et al.*, 2006, 2007). While technical difficulties and high cost can make it difficult to analyze a large number of samples by the clone library method, cataloguing bacterial species in the oral cavity by clone libraries provides 16S rRNA gene sequence data that can be used to design primers or oligonucleotide probes to target both cultivable and as-yet-uncultivated bacteria in infected root canals. Primers and probes can be used in PCR and DNA-DNA hybridization assays (*e.g.*, reverse-capture

checkerboard, microarrays) in large-scale clinical studies to investigate prevalence and association with disease (Rôças and Siqueira, 2005b, 2008; Siqueira and Rôças, 2005c, 2009a).


## DIVERSITY OF THE ENDODONTIC MICROBIOTA—OVERALL FINDINGS

'Microbiota' is arguably the best collective term for microorganisms, since widely used terms such as 'microbial flora' and 'microflora' perpetuate an outdated classification of microorganisms as plants (Dethlefsen *et al.*, 2006). 'Diversity' is a function of both the number of species present in a community (species richness) and the relative abundance of those species in the system (species evenness). The highest diversity occurs in communities with many different species present (richness) in relatively equal abundance (evenness) (Huston, 1994). The richness and evenness of microbial communities are the result of selective pressures that shape diversity within communities. This review discusses the diversity of the endodontic microbiota, with the main focus on the aspect of species richness or composition. The integration of data from culture-dependent and culture-independent methods for microbial identification in endodontic infections is presented here for the first time.

Representatives of the domains *Archaea* and *Eukarya* have been occasionally reported to occur in endodontic infections. *Archaea* diversity is restricted to an as-yet-uncultivated *Methanobravibacter oralis*-like phylotype (Vianna *et al.*, 2006a; Vickerman *et al.*, 2007). Except for a couple of molecular studies targeting *Candida albicans* (Baumgartner *et al.*, 2000; Siqueira and Rôças, 2004a), the diversity of *Eukarya* in infected canals has been assessed exclusively by culture-dependent methods. Six *Candida*, one *Geotrichum*, one *Rhodotorula*, and one *Saccharomyces* species have been isolated from root canals (see Appendix for references).

Although a few representatives of the domains *Eukarya* and *Archaea* have been found in endodontic infections, and reports indicate the occurrence of herpesviruses (cytomegalovirus and Epstein-Barr virus) in apical periodontitis lesions (Sabeti *et al.*, 2003), the domain *Bacteria* is far more dominant and diverse in endodontic infections. Integrated datasets from culture and molecular studies published as of March 2008, showed that 468 unique bacterial taxa have been found in endodontic infections associated with different clinical conditions. The vast majority of these bacterial taxa fall into 100 genera, while 22 taxa remained undefined and were assigned to family or even to phylum level. *Prevotella* (39 taxa), *Eubacterium* (27 taxa), *Streptococcus* (26 taxa), and *Lactobacillus* (21 taxa) are the most represented genera. All taxa already reported in endodontic infections and the respective studies that detected them are listed in the Appendix.

At the broader phylogenetic level, all the detected bacterial taxa have been found to belong to 9 of the 13 phyla that have oral representatives, namely, *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, *Fusobacteria*, *Spirochaetes*, *Synergistes*, TM7, and SR1 (Table 1). The most common representatives from each of these phyla are shown in Fig. 1. Members of *Acidobacteria*, a phylum sporadically found in the oral cavity, were not detected by group-specific PCR (unpublished data from the authors' laboratory).



**Figure 1.** Bacterial phyla that have representatives in endodontic infections. On the right, example species or phylotypes for each phylum are presented.

It is important to point out that the numbers of bacterial taxa reported refer exclusively to richness within each phylum and have no relationship to prevalence and importance in disease causation. For instance, the phyla *Fusobacteria*, *Spirochaetes*, and *Synergistes* have few representative species/phylotypes in endodontic infections, but some of them, including *Fusobacterium nucleatum*, *Treponema denticola*, and *Synergistes* clone BA121, respectively, are among the most commonly detected taxa in primary intraradicular infections.

Of the recovered taxa, 204 (44%) were isolated or detected in only one study. This may indicate that they are subject-specific and then contribute to the high level of interindividual variation in bacterial community profiles, or they may be contaminants or species poorly identified. Contaminants may

be introduced in the clinical setting during sample collection (usually oral contaminants) or in the laboratory during specimen handling (non-oral contaminants). Also, “single-study” taxa were rarely very prevalent in the individual study that reported their occurrence. In contrast, as-yet-uncultivated phylotypes or culture-difficult species detected in only one molecular study may not necessarily be interpreted as irrelevant because their occurrence in a single study may reflect the fact that, thus far, only a few broad-range molecular studies have been published, and the microbiota have been investigated in only a limited number of individuals.

The analysis of the bacteria detected in endodontic infections revealed that 210 distinct bacterial taxa were reported in molecular biology studies alone. This figure corresponds to 45% of all taxa already found in endodontic infections, as opposed to 151 taxa (32%) detected by culture studies alone. One hundred seven taxa, 23% of the total bacterial species richness, were detected by the application of both culture-dependent and culture-independent techniques (Fig. 2).

Culture-dependent and culture-independent molecular studies provide a somewhat different description of the diversity of the endodontic microbiota, as indicated by the degree of overlapping findings (Fig. 2). However, both approaches show unequivocally that *Firmicutes* are by far the most diverse group, even though the community structure of this group has been largely underestimated in the reports from culture-dependent studies. Representatives of 4 phyla—*Spirochaetes*, *Synergistes*, TM7, and SR1—have been found almost exclusively in endodontic infections by molecular

**Table 1.** Overall Findings of Bacterial Species/Phylotype (taxa) Richness in the Different Types of Endodontic Infections

Phyla	Taxa	As-Yet-	Taxa Detected	Taxa Detected
		Uncultivated	by Molecular	by Culture
		Phylotypes	Studies	Studies
<i>Firmicutes</i>	220	81	146	124
<i>Bacteroidetes</i>	73	28	47	48
<i>Actinobacteria</i>	69	21	43	42
<i>Proteobacteria</i>	65	17	44	34
<i>Fusobacteria</i>	15	6	11	9
<i>Spirochaetes</i>	14	4	14	0
<i>Synergistes</i>	10	10	10	1
TM7	1	1	1	0
SR1	1	1	1	0


biology methods. While the latter two have only one representative each, both of which have been detected in low prevalence, members of the two former have been as frequent as, or even more frequent than, most representatives of the other, more diverse, phyla.

Molecular studies have allowed for the detection of 169 phylotypes representing bacteria that have not yet been cultivated and/or fully characterized. Most of these phylotypes belong to the phyla *Firmicutes*, *Bacteroidetes*, and *Actinobacteria*. As-yet-uncultivated and/or uncharacterized phylotypes can be important pathogens that have been overlooked by culture-dependent studies.

Molecular methods have succeeded over culture methods in the detection of not only as-yet-uncultivated and uncharacterized phylotypes, but also of cultivable and validly named species, since about 40 taxa included in this category have been exclusively detected by molecular approaches. These bacteria either represent fastidious species that were not recovered (probably because specific culture media were not used), or they are species that were successfully cultivated but inadequately identified, because methods improper for definite identification were used or the isolate presented ambiguous phenotypic behavior (Siqueira *et al.*, 2007c). Indeed, application of the 16S rRNA gene-sequencing approach for the identification of culture isolates has demonstrated that some phylotypes previously supposed to be “uncultivable” can actually be cultivated by conventional anaerobic techniques with ordinary growth media (Munson *et al.*, 2002; Siqueira *et al.*, 2007c). Most of these previously “uncultivable” bacteria remain uncharacterized, and a valid species name is pending.

The strength of molecular methods to unravel the bacterial diversity and its superiority over culture approaches in this regard is in clear agreement with studies in other areas (Amann *et al.*, 1995; Rappe and Giovannoni, 2003; Janssen, 2006; Rajilić-Stojanović *et al.*, 2007). In endodontic microbiology research, it is quite remarkable that molecular studies performed over the past 10 years have revealed 317 distinct bacterial taxa in infected canals, while culture studies over the past 30 years revealed 258 taxa.

Bacterial taxa for which there is moderate-to-strong evidence of involvement with causation of apical periodontitis are shown in Fig. 3. Selection of these taxa was based on their detection in several independent studies and moderate-to-high prevalence in more than one study. Thus, the taxa shown in Fig. 3 can be regarded as the main candidate endodontic pathogens associated with different types of infection. One will notice that there are some ubiquitous endodontic taxa that have been found by at least one study in every type of endodontic infection. Most of these “cosmopolitan” species have been found to be very prevalent in association with some specific



**Figure 2.** Distribution of bacterial species/phylotypes found in endodontic infections according to the detection method. Data are given overall and for the 9 phyla that have endodontic representatives.


types of infection, mainly primary infections. In spite of having been found in samples from every type of infection, *Enterococcus faecalis* is encountered more frequently in treated root canals of teeth evincing post-treatment disease (Appendix). All of these ubiquitous bacteria are cultivable and validly named species. However, it is worth pointing out that this list is biased by the fact that, for some types of infection, there is no or only one molecular study using broad-range analysis. Thus, some taxa found in primary infections in even high prevalence may have been deprived of this “status” because of the lack of a body of molecular studies investigating other disease categories.

## PRIMARY INTRARADICULAR INFECTIONS

Primary intraradicular infection is caused by micro-organisms that initially invade and colonize the necrotic pulp tissue. It is characterized by a mixed consortium conspicuously dominated by anaerobic bacteria and composed of 10 to 30 species *per* canal (Munson *et al.*, 2002; Siqueira *et al.*, 2004b; Siqueira and Rôças, 2005b). Total bacterial counts vary from  $10^3$  to  $10^8$  cells *per* infected canal (Sundqvist, 1976; Vianna *et al.*, 2006b; Sakamoto *et al.*, 2007; Siqueira *et al.*, 2007d).

Bacterial profiles of the endodontic microbiota vary from individual to individual (Siqueira *et al.*, 2004b; Sakamoto *et al.*, 2006), *i.e.*, each individual harbors a unique endodontic microbiota in terms of species richness and abundance. This indicates that primary apical periodontitis has a heterogeneous etiology, where no single species can be considered as the main endodontic pathogen, and multiple bacterial combinations play a role in disease causation. This is consistent with the high numbers of different taxa that have been detected in primary infections.

Collectively, culture and molecular methods have allowed 391 bacterial, 4 fungal, and one archaeal taxa to be detected in primary



**Figure 3.** Phylogenetic tree based on 16S rRNA gene comparisons, showing several candidate endodontic pathogens and the clinical conditions with which they have been associated. Distribution of bacterial species/phylotypes among the different types of infections is shown by the columns of boxes to the right of the tree. Different darknesses of the box indicate strength of association based on prevalence data and the number of studies in which the species was detected, *i.e.*, clear box means not detected, while black box means detected by several studies and/or in high prevalence. The reference bar indicates 5% sequence divergence. *Pi*, primary infections; *Fs*, filling stage; *Rc*, retreatment cases; *Ex*, extraradicular infections.

infections. As-yet-uncultivated and/or uncharacterized bacteria have been represented by 136 taxa. Molecular methods have identified 271 taxa, while 216 taxa have been isolated by culture.

Bacterial species/phylotypes detected in primary infections fall into the 9 phyla mentioned above and 82 genera. Seventeen taxa have not been assigned to a genus, only to family or phylum. The highest species richness was observed for the *Firmicutes*, followed by *Bacteroidetes* and *Actinobacteria* (Table 2). Of the 391 taxa detected, 261 were unique to primary infections, *i.e.*, they have not been detected in samples from other types of infections.

Diverse groups of Gram-negative and Gram-positive bacteria have been identified in primary infections. Black-pigmented Gram-negative anaerobic rods are named after their ability to form brown to black colonies on blood agar plates and have been classified into two genera—*Prevotella* (containing saccharolytic species) and *Porphyromonas* (containing asaccharolytic species). The genus *Prevotella* also includes some bile-sensitive non-pigmented species. *Prevotella* species, especially *P. intermedia*, *P. nigrescens*, *P. tanneriae*, *P. baroniae*, and *P. denticola*, have been among the most frequently detected taxa in primary infections (see Appendix for references). Of the *Porphyromonas* species, *P. endodontalis* and *P. gingivalis* have been consistently encountered in endodontic infections, and a role in the etiology of acute abscesses is suspected (van Winkelhoff *et al.*, 1985; Haapasalo *et al.*, 1986; Sundqvist *et al.*, 1989; Siqueira *et al.*, 2001a; Seol *et al.*, 2006).

An important periodontal pathogen, *Tannerella forsythia* (formerly *Bacteroides forsythus*), a fastidious Gram-negative obligate anaerobe that had never been detected in root canals by culture, was for the first time reported to occur in primary endodontic infections in a study with species-specific PCR (Conrads *et al.*, 1997). Subsequent studies with different molecular biology approaches have confirmed that *T. forsythia* is a common member of the microbiota

associated with different types of primary infections, including abscesses (Jung *et al.*, 2000; Siqueira *et al.*, 2000b, 2001b; Fouad *et al.*, 2002; Siqueira and Rôças, 2003c).

*Dialister* species are asaccharolytic obligately anaerobic Gram-negative coccobacilli that represent another example of bacteria that have been consistently detected in endodontic infections only after the advent of molecular biology techniques. *D. pneumosintes* and the recently described *D. invisus* have been frequently present in the microbiota associated with asymptomatic and symptomatic primary infections (Appendix).

*Fusobacterium nucleatum* is an anaerobic spindle-shaped rod that is one of the most commonly encountered Gram-negative species in endodontic infections. *F. nucleatum* has been frequently isolated from or detected in primarily infected root canals as well as in endodontic abscesses (Appendix).

Spirochetes are highly motile spiral-shaped bacteria that have been frequently observed by microscopy in samples taken from endodontic infections. Nonetheless, they had never been identified to the species level. The application of molecular diagnostic methods to the identification of spirochetes has demonstrated that occurrence of these spiral bacteria in endodontic infections has been overlooked by technical hurdles of culture techniques. All of the currently cultivable and named oral *Treponema* species have been disclosed by molecular studies of primary endodontic infections (Appendix). The most predominant species in primary infections are *T. denticola* and *T. socranskii* (Siqueira *et al.*, 2000b; Baumgartner *et al.*, 2003; Rôças *et al.*, 2003; Siqueira and Rôças, 2004b).

Even though anaerobic Gram-negative bacteria have been found to be the most common micro-organisms in primary infections, several Gram-positive rods have also been frequent members of the endodontic microbial consortium. Of these, *Pseudoramibacter alactolyticus* has been frequently isolated from or detected in samples from endodontic infections, in prevalence values as high as those of the most commonly found Gram-negative species (Sundqvist, 1992; Siqueira and Rôças, 2003d; Siqueira *et al.*, 2004a). *Filifactor alocis* is an obligately anaerobic rod that has been detected by molecular studies in about one-half of the samples from primary infections (Siqueira and Rôças, 2003a; Gomes *et al.*, 2006). *Actinomyces* species, which have been associated with failure of the endodontic treatment by causing apical actinomycosis, have been found in about 10% of infected canals (Sundqvist, 1992; Siqueira *et al.*, 2002). *Propionibacterium propionicum*, another species that can participate in apical actinomycosis, has also been commonly detected in samples from primary infections (Siqueira and Rôças, 2003e). *Olsenella* species consist of small non-motile Gram-positive obligately anaerobic rods, which represent another example of bacteria that have been detected in endodontic infections only after the introduction of molecular biology methods (Fouad *et al.*, 2002; Munson *et al.*, 2002). Among the *Olsenella* clade, *O. uli* has been the most commonly found species in endodontic infections (Chávez de Paz *et al.*, 2004; Rôças and Siqueira, 2005a).

Gram-positive cocci, specifically peptostreptococci and streptococci, are frequently present in primary endodontic infections. *Parvimonas* (formerly *Peptostreptococcus* or *Micromonas*) *micra* is an asaccharolytic anaerobic Gram-positive small coccus that has

**Table 2.** Bacterial Species/Phylotype (taxa) Richness in Primary Endodontic Infections

Phyla	Taxa	As-Yet-Uncultivated Phylotypes	Taxa Detected by Molecular Studies	Taxa Detected by Culture Studies
<i>Firmicutes</i>	184	69	131	98
<i>Bacteroidetes</i>	69	24	42	48
<i>Actinobacteria</i>	54	11	31	39
<i>Proteobacteria</i>	44	11	32	21
<i>Fusobacteria</i>	14	5	9	9
<i>Spirochaetes</i>	14	4	14	0
<i>Synergistes</i>	10	10	10	1
TM7	1	1	1	0
SR1	1	1	1	0

been isolated from or detected in about one-third of the primarily infected canals, and its prevalence in symptomatic infections has also been relatively high (Sundqvist, 1992; Weiger *et al.*, 1995; Gomes *et al.*, 1996; Khemalaelakul *et al.*, 2002; Siqueira *et al.*, 2003; Chu *et al.*, 2005). Members of the *Streptococcus anginosus* group have been reported to be the most prevalent streptococci, but *S. oralis*, *S. mitis*, and *S. sanguinis* can also often be recovered/detected (Sundqvist, 1992; Siqueira *et al.*, 2002).

*Campylobacter* species, including *C. rectus* and *C. gracilis*, are Gram-negative anaerobic rods that have been detected in primary endodontic infections, but in low-to-moderate prevalence values (Sundqvist, 1992; Le Goff *et al.*, 1997; Siqueira and Rôças, 2003b). *Catonella morbi*, a saccharolytic obligately anaerobic Gram-negative rod associated with marginal periodontitis, has been found in about one-fourth of the cases of primary endodontic infections (Siqueira and Rôças, 2006). Other bacteria detected in low-to-moderate frequencies in primary infections include: *Veillonella parvula*, *Eikenella corrodens*, *Neisseria mucosa*, *Centipeda periodontii*, *Granulicatella adiacens*, *Gemella morbillorum*, *Capnocytophaga* species, and anaerobic lactobacilli (Appendix).

Investigations of the diversity of the microbiota of primary infections by broad-range PCR and 16S rRNA gene clone library analysis have demonstrated that it is far more complex than previously reported by culture studies. Noteworthy is the common occurrence of as-yet-uncultivated bacteria—about 40-55% of the endodontic microbiota is composed of as-yet-uncultivated phylotypes (Munson *et al.*, 2002; Sakamoto *et al.*, 2006, 2007). A molecular study investigating primary infections revealed that 55% and 40% of the taxa detected in association with chronic apical periodontitis and acute apical abscesses were as-yet-uncultivated phylotypes, respectively (Sakamoto *et al.*, 2006). As for their abundance in these infections, as-yet-uncultivated phylotypes corresponded to 38% and 30% of the clones sequenced from samples of chronic apical periodontitis and abscesses, respectively (Sakamoto *et al.*, 2006).

As mentioned above, uncultivated phylotypes have been reported for all phyla with endodontic representatives. Several of these phylotypes can be candidate endodontic pathogens based on association data. For instance, oral *Synergistes* clones BA121, E3\_33, BH017, and W090, which had been originally assigned to the *Flexistipes* or *Deferribacteres* group (Paster *et al.*, 2001; Godon

*et al.*, 2005), have been commonly detected in samples from asymptomatic and symptomatic endodontic infections (Rôças and Siqueira, 2005b; Siqueira and Rôças, 2005c; Siqueira *et al.*, 2005; Vianna *et al.*, 2007). The great majority of *Synergistes* bacteria remain uncultivated, and this can be the primary reason for the fact that their presence in endodontic infections has been overlooked by culture studies. Similarly, clones I025 and X112 from the TM7 and SR1 candidate phyla, respectively, which so far have no cultivable representatives, have also been detected in primary infections (Rôças and Siqueira, 2005a, 2008; Siqueira and Rôças, 2005c). Uncultivated phylotypes related to several genera have been disclosed, mainly *Dialister*, *Prevotella*, *Peptostreptococcus*, *Solobacterium*, *Olsenella*, *Selenomonas*, *Eubacterium*, *Megasphaera*, and *Veillonella*, as well as phylotypes related to the family *Lachnospiraceae* (Rolph *et al.*, 2001; Munson *et al.*, 2002; Saito *et al.*, 2006; Sakamoto *et al.*, 2006; Vickerman *et al.*, 2007). One study (Sakamoto *et al.*, 2006) found some uncultivated phylotypes among the most prevalent bacteria in primary infections, including *Lachnospiraceae* oral clone 55A-34, *Megasphaera* oral clone CS025, and *Veillonella* oral clone BP1-85. *Prevotella* oral clone PUS9.180, *Eubacterium* oral clone BP1-89, and *Lachnospiraceae* oral clone MCE7\_60 were exclusively detected in symptomatic samples (Sakamoto *et al.*, 2006). Uncultivated phylotypes are previously unrecognized and overlooked bacteria that may play a role in the pathogenesis of different forms of apical periodontitis.

## PERSISTENT/SECONDARY INTRARADICULAR INFECTIONS

Secondary infection is caused by micro-organisms that were not present in the primary infection, but were introduced in the root canal at some time after professional intervention (so called because it is secondary to intervention) (Siqueira, 2002). Persistent infection is caused by micro-organisms that were members of a primary or secondary infection and that, in some way, resisted intracanal antimicrobial procedures and endured periods of nutrient deprivation in treated canals (Siqueira, 2002).

Persistent or secondary intraradicular infections are the major causes of several clinical problems, including endodontic treatment failure, which is characterized by persistence or appearance of apical periodontitis after treatment (Siqueira, 2008). Studies have attempted to identify the micro-organisms found at the root-canal-filling stage, which may jeopardize the treatment outcome, and the micro-organisms in root-canal-treated teeth with apical periodontitis, which are arguably the cause of failure.

Analysis of integrated data from culture and molecular studies revealed that 51 bacterial and 2 fungal taxa have been detected in samples taken at the time of filling and also at the time of re-treatment. The following taxa have been detected by several studies: *Propionibacterium acnes*, *P. propionicum*, *Actinomyces naeslundii*, *Actinomyces odontolyticus*, *P. intermedia*, *Anaerococcus prevotii*, *Eggerthella lenta*, *E. faecalis*, *G. morbillorum*, *P. micra*, *P. alactolyticus*, *S. anginosus* group, *S. mitis*, *F. nucleatum*, and *C. albicans*. Theoretically, taxa found at both the time of filling and during re-treatment may be involved in persistent infections. Likewise, taxa that were found only at the time of re-treatment, but not at the time of filling, may represent

secondary infections that developed by lack of a bacteria-tight coronal seal. Taxa detected at the filling stage, but not at the time of re-treatment, may not be able to endure the conditions within obturated canals. Although all this discussion sounds logical and interesting, it is largely speculative, because the data belong to separate cross-sectional studies, and no strong evidence can be taken in this regard.

## BACTERIA AT THE ROOT-CANAL-FILLING STAGE (POST-TREATMENT SAMPLES)

Root canal samples can be taken at the time of treatment to assess the antimicrobial efficacy of treatment protocols and/or check the bacteriologic conditions of the canal before filling procedures. Samples positive for bacterial growth after chemomechanical preparation, followed (or not) by intracanal medication, have been shown to harbor an average of 1 to 5 bacterial species, at counts reaching up to  $10^2$  to  $10^5$  cells *per* canal (Byström and Sundqvist, 1985; Sjögren *et al.*, 1997; Vianna *et al.*, 2006b; Sakamoto *et al.*, 2007; Siqueira *et al.*, 2007a,b). These numbers indicate that, even if total bacterial elimination does not occur, at least a substantial reduction in species diversity is attained after treatment.

Culture and molecular methods have allowed 103 bacterial and 6 fungal taxa to be detected in samples taken after chemomechanical preparation and/or intracanal medication. Molecular methods have detected 26 taxa, while 88 taxa were isolated in culture studies. Forty-one taxa were found in only one study.

Bacterial species/phylotypes detected in post-treatment samples belong to 5 phyla and 41 genera. Three taxa have been assigned to taxonomic levels above genus. The highest species richness was observed for the *Firmicutes*, followed by *Proteobacteria* and *Actinobacteria* (Table 3).

Gram-negative bacteria, which are common members of primary infections, are usually eliminated after treatment procedures (Chávez de Paz, 2005). Exceptions may include some anaerobic rods, such as *F. nucleatum*, *Prevotella* species, and *C. rectus*, which are among the species found in post-instrumentation/medication samples (Appendix). However, most studies have clearly revealed that, when bacteria resist treatment procedures, Gram-positive bacteria are more frequently present. They include streptococci (*S. mitis*, *S. gordonii*, *S. anginosus*, *S. sanguinis*, and *S. oralis*), *P. micra*, *Actinomyces* species (*A. israelii* and *A. odontolyticus*), *Propionibacterium* species (*P. acnes* and *P. propionicum*), *P. alactolyticus*, lactobacilli (*L. paracasei* and *L. acidophilus*), *E. faecalis*, and *O. uli* (Appendix). Other Gram-positive bacteria, including *Bifidobacterium* species, *Eubacterium* species, and staphylococci, can also be found, but in lower frequencies (Sjögren *et al.*, 1997; Chávez de Paz, 2004; Chu *et al.*, 2006).

With the recent findings showing as-yet-uncultivated bacteria as constituents of a significant proportion of the endodontic microbiota (Munson *et al.*, 2002; Sakamoto *et al.*, 2006), studies on the effects of intracanal antimicrobial procedures should also focus on these bacteria. Thus far, the only broad-range molecular study to investigate samples from this condition used a 16S rRNA gene clone library analysis to identify the bacteria persisting after chemomechanical preparation, with 2.5% NaOCl as irrigant and intracanal medication with a calcium hydroxide paste (Sakamoto

**Table 3.** Bacterial Species/Phylotype (taxa) Richness in Samples Taken at the Time of Root Canal Filling

Phyla	Taxa	As-Yet- Uncultivated Phylotypes	Taxa Detected by Molecular Studies	Taxa Detected by Culture Studies
<i>Firmicutes</i>	45	2	10	40
<i>Proteobacteria</i>	20	3	3	17
<i>Actinobacteria</i>	19	2	9	15
<i>Bacteroidetes</i>	16	3	3	13
<i>Fusobacteria</i>	3	0	1	3

*et al.*, 2007). Eleven, 4, and 5 taxa were detected in initial (S1), post-instrumentation (S2), and post-medication (S3) samples, respectively. No taxon found in post-treatment samples was shown to dominate the initial (primary) infection (baseline). *Streptococcus* species were detected in all post-treatment samples and were also the most dominant taxa in these samples, except for a S2 sample in which *Solobacterium* clone K010 corresponded to 56% of the clones sequenced. Forty-two percent of the taxa found in post-treatment samples were as-yet-uncultivated bacteria, indicating that previously uncharacterized bacteria may also persist after treatment, and their role in the long-term treatment outcome requires elucidation.

## MICROBIOTA IN ROOT-CANAL-TREATED TEETH (RE-TREATMENT SAMPLES)

Root canal samples can be taken from teeth that were treated months to years previously and need endodontic re-treatment because of the emergence or persistence of disease. The microbiota in these cases also exhibit a decreased diversity in comparison with that of primary infections. Canals apparently well-treated contain from 1 to 5 species, while the number of species in canals with inadequate treatment can reach 30, which is very similar to that in untreated canals (Sundqvist *et al.*, 1998; Pinheiro *et al.*, 2003; Rôças *et al.*, 2004c; Siqueira and Rôças, 2004a). A single treated canal associated with post-treatment disease can harbor a density of  $10^3$  to  $10^7$  bacterial cells (Peciulienė *et al.*, 2001; Sedgley *et al.*, 2006).

Integrated datasets from culture and molecular methods have revealed 158 bacterial and 3 fungal taxa in samples taken at the time of root canal re-treatment of teeth evincing apical periodontitis lesions. Forty-six as-yet-uncultivated phylotypes have been detected by molecular studies. Molecular methods have detected 109 taxa, while culture studies isolated 72 taxa. A great many taxa (106) have been reported by only one study.

Bacterial species/phylotypes detected in re-treatment samples belong to 7 phyla and 58 genera. Nine taxa have been assigned to the family or phylum level. The highest species richness was again observed for the *Firmicutes*, followed by *Actinobacteria* and *Proteobacteria* (Table 4). Molecular studies have allowed members of the *Spirochaetes* and *Synergistes* phyla to be detected in these cases.

Culture-dependent and culture-independent studies have revealed that *E. faecalis* is the most frequent species in root-canal-treated teeth, with prevalence values reaching up to 90% of the cases (Molander *et al.*, 1998; Sundqvist *et al.*, 1998; Pinheiro *et al.*, 2003;

**Table 4.** Bacterial Species/Phylotype (taxa) Richness in Persistent/Secondary Infections Related to Treatment Failure

Phyla	Taxa	As-Yet- Uncultivated Phylotypes	Taxa Detected by Molecular Studies	Taxa Detected by Culture Studies
<i>Firmicutes</i>	76	21	50	40
<i>Actinobacteria</i>	28	11	21	10
<i>Proteobacteria</i>	26	5	17	11
<i>Bacteroidetes</i>	22	6	16	9
<i>Fusobacteria</i>	4	2	3	2
<i>Spirochaetes</i>	1	0	1	0
<i>Synergistes</i>	1	1	1	0

Rôças *et al.*, 2004a,b; Siqueira and Rôças, 2004a; Sedgley *et al.*, 2006; Zoletti *et al.*, 2006). Root-canal-treated teeth are about 9 times more likely to harbor *E. faecalis* than cases of primary infections (Rôças *et al.*, 2004a). Other bacteria found in re-treatment cases include streptococci and some fastidious anaerobic bacterial species—*P. alactolyticus*, *P. propionicum*, *F. alocis*, *D. pneumosintes*, and *D. invisus* (see Appendix).

In a study using 16S rRNA gene clone library analysis, Sakamoto *et al.* (2008) found a mean of 10 taxa per treated root canal, ranging from 1 to 26. As-yet-uncultivated phylotypes corresponded to 55% of the taxa detected. Only 11 taxa were found in more than one case: *Bacteroidetes* oral clone X083, an uncultured *Saprosiraceae* clone, *Prevotella oris*, *P. baroniae*, *Pseudomonas aeruginosa*, *Streptococcus mutans*, *Synergistes* clone BA121, *Dialister* clone 9N-1, *E. faecalis*, *Flavobacteriaceae* genomospecies C1, and *Peptostreptococcus* clone FG014. A high interindividual variability in the composition of the microbiota was clearly evident, and, along with the findings from a bacterial community profiling technique (Rôças *et al.*, 2004c), suggests that distinct bacterial combinations can play a role in treatment failures. Bacterial abundance also varied from sample to sample, and some uncultivated phylotypes were the most dominant taxa in about one-half of the cases examined. In fact, uncultivated phylotypes made up a significant fraction of the microbiota in root-canal-treated teeth, since they comprised about 50% of the total number of clones sequenced (Sakamoto *et al.*, 2008).

Fungi are only occasionally found in primary infections, but *Candida* species have been detected in root-canal-treated teeth in up to 18% of the cases (Cheung and Ho, 2001; Peciulienė *et al.*, 2001; Egan *et al.*, 2002). *C. albicans* is by far the most commonly detected fungal species in re-treatment cases.

## EXTRARADICULAR INFECTIONS

Extraradicular infection is characterized by microbial invasion of and proliferation in the inflamed periradicular tissues, and is derived from intraradicular infection (Tronstad *et al.*, 1987; Siqueira, 2002; Tronstad and Sunde, 2003). Extraradicular infection in cases of chronic apical periodontitis represents one of the greatest controversies in endodontics (Bergenholtz and Spangberg, 2004). The infection is conceivably limited to a small proportion of cases, and, when present, it can be dependent on or independent of intraradicular infection (Siqueira,

**Table 5.** Bacterial Species/Phylotype (taxa) Richness in Extraradicular Infections

Phyla	Taxa	Taxa Detected by Molecular Studies	Taxa Detected by Culture Studies
<i>Firmicutes</i>	47	16	41
<i>Bacteroidetes</i>	18	10	12
<i>Proteobacteria</i>	12	6	8
<i>Actinobacteria</i>	10	7	8
<i>Fusobacteria</i>	3	3	2
<i>Spirochaetes</i>	2	2	0

2008). A study revealed that the number of species *per* infected lesion ranged from 11 to 34 (Sunde *et al.*, 2000).

Culture and molecular methods have revealed 92 bacterial taxa and one fungal species in samples taken from apical periodontitis lesions associated with treated canals. No reliable broad-range molecular study has thus far been conducted to survey these infections. For instance, results from samples taken from extracted teeth were not considered herein because of the obvious risks of contamination of the lesion. Therefore, molecular methods to investigate the bacteria in extraradicular infections associated with treated teeth evaluated only cultivable taxa. These molecular methods alone have detected 44 taxa, while 71 species have been isolated in culture studies. Forty-six taxa were found in a single study.

Bacterial taxa detected in extraradicular infections belong to 6 phyla and 38 genera. The *Firmicutes* was the phylum with the largest number of representatives, followed by *Bacteroidetes* and *Proteobacteria* (Table 5). Species reported by many studies include: *Actinomyces* species (*A. israelii*, *A. naeslundii*, *A. odontolyticus*, *A. viscosus*), *P. acnes*, *P. propionicum*, *P. gingivalis*, *P. intermedia*, *Prevotella oralis*, *P. micra*, and *F. nucleatum*.

## CONCLUDING REMARKS

The integration of datasets from culture-dependent and culture-independent molecular biology studies has shown that 468 unique bacterial, one archaeal, and 9 fungal taxa have been, as of then, reported to occur in endodontic infections associated with different clinical conditions. Of the bacterial taxa, 264 have been found in more than one study.

It is important to point out that these figures might be somewhat reduced had only studies with stringent culture (*e.g.*, biochemical tests and gas-liquid chromatography) and/or molecular (*e.g.*, 16S rRNA gene sequencing) identification approaches been surveyed. However, this would have significantly reduced the number of studies evaluated, with the inevitable risk of excluding many important species that can be easily identified by less stringent approaches. Moreover, several other factors may have influenced the number of species/phylogenotypes compiled in this study, including the protocol for disinfection of the operative field, as well as the care, level of training, and experience during sample collection, sample handling in the laboratory, and interpretation of identification data. All of these factors may have influenced the species/phylogenotype numbers in our list. However, we could not remediate this by ruling out some studies, since most of these factors cannot be measured based on a printed paper.

While there was an apparent difference in the prevalence of some of the bacterial species found in primary infection *vs.* persistent or secondary infections, many species were present in only one specific study but not in others, making it difficult to determine their association with any specific form of endodontic infection. Several of these species are culture-difficult fastidious species, newly named taxa, or as-yet-uncultivated phylogenotypes, for which studies are very recent, with current molecular technology, and no further corroborative findings have become available. Further prevalence studies should target such species, especially those very prevalent in a single isolated study, to check their actual prevalence and look for associations with different forms of disease.

Now that a comprehensive inventory of the microbial species present in these infections has been established by culture and molecular studies, further efforts should be directed toward finding associations of the most prevalent taxa with symptoms and other clinical conditions, unraveling the functional role of the detected species in the mixed endodontic consortium, and disclosing their susceptibility to antimicrobial substances and treatment procedures. Extensive review of the endodontic microbiology literature leads to the conclusion that apical periodontitis has a polymicrobial etiology, and that the different types of endodontic infections are usually represented by a mixed consortium, whose diversity varies according to the type of infection. The bacterial community profile exhibits a high interindividual variability, and differences are even more pronounced when samples are taken from different geographical locations.

## ACKNOWLEDGMENT

This work is supported by grants from CNPq, a Brazilian Governmental Institution.

## REFERENCES

- Aas JA, Paster BJ, Stokes LN, Olsen I, Dewhirst FE (2005). Defining the normal bacterial flora of the oral cavity. *J Clin Microbiol* 43:5721-5732.
- Aas JA, Barbuto SM, Alpagot T, Olsen I, Dewhirst FE, Paster BJ (2007). Subgingival plaque microbiota in HIV positive patients. *J Clin Periodontol* 34:189-195.
- Amann RI, Ludwig W, Schleifer KH (1995). Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiol Rev* 59:143-169.
- Baumgartner JC, Falkler WA Jr (1991). Bacteria in the apical 5 mm of infected root canals. *J Endod* 17:380-383.
- Baumgartner JC, Watts CM, Xia T (2000). Occurrence of *Candida albicans* in infections of endodontic origin. *J Endod* 26:695-698.
- Baumgartner JC, Khemaleelakul SU, Xia T (2003). Identification of spirochetes (treponemes) in endodontic infections. *J Endod* 29:794-797.
- Bergenholtz G (1974). Micro-organisms from necrotic pulp of traumatized teeth. *Odontol Revy* 25:347-358.
- Bergenholtz G, Spangberg L (2004). Controversies in endodontics. *Crit Rev Oral Biol Med* 15:99-114.
- Bik EM, Eckburg PB, Gill SR, Purdom EA, Francois F, *et al.* (2006). Molecular analysis of the bacterial microbiota in the human stomach. *Proc Natl Acad Sci USA* 103:732-737.
- Byström A, Sundqvist G (1985). The antibacterial action of sodium hypochlorite and EDTA in 60 cases of endodontic therapy. *Int Endod J* 18:35-40.
- Chávez de Paz L (2004). Gram-positive organisms in endodontic infections. *Endod Topics* 9:79-96.
- Chávez de Paz LE (2005). On bacteria persisting root canal treatment. Identification and potential mechanisms of resistance to antimicrobial measures [PhD thesis]. Göteborg, Sweden: Göteborg University.

- Chávez de Paz LE, Molander A, Dahlén G (2004). Gram-positive rods prevailing in teeth with apical periodontitis undergoing root canal treatment. *Int Endod J* 37:579-587.
- Cheung GS, Ho MW (2001). Microbial flora of root canal-treated teeth associated with asymptomatic periapical radiolucent lesions. *Oral Microbiol Immunol* 16:332-337.
- Chu FC, Tsang CS, Chow TW, Samaranyake LP (2005). Identification of cultivable microorganisms from primary endodontic infections with exposed and unexposed pulp space. *J Endod* 31:424-429.
- Chu FC, Leung WK, Tsang PC, Chow TW, Samaranyake LP (2006). Identification of cultivable microorganisms from root canals with apical periodontitis following two-visit endodontic treatment with antibiotics/steroid or calcium hydroxide dressings. *J Endod* 32:17-23.
- Conrads G, Gharbia SE, Gulabivala K, Lampert F, Shah HN (1997). The use of a 16S rDNA directed PCR for the detection of endodontopathogenic bacteria. *J Endod* 23:433-438.
- Dekio I, Hayashi H, Sakamoto M, Kitahara M, Nishikawa T, Suematsu M, et al. (2005). Detection of potentially novel bacterial components of the human skin microbiota using culture-independent molecular profiling. *J Med Microbiol* 54(Pt 12):1231-1238.
- de Lillo A, Ashley FP, Palmer RM, Munson MA, Kyriacou L, Weightman AJ, et al. (2006). Novel subgingival bacterial phylotypes detected using multiple universal polymerase chain reaction primer sets. *Oral Microbiol Immunol* 21:61-68.
- Dethlefsen L, Eckburg PB, Bik EM, Relman DA (2006). Assembly of the human intestinal microbiota. *Trends Ecol Evol* 21:517-523.
- Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, et al. (2005). Diversity of the human intestinal microbial flora. *Science* 308:1635-1638.
- Egan MW, Spratt DA, Ng YL, Lam JM, Moles DR, Gulabivala K (2002). Prevalence of yeasts in saliva and root canals of teeth associated with apical periodontitis. *Int Endod J* 35:321-329.
- Fouad AF, Barry J, Caimano M, Clawson M, Zhu Q, Carver R, et al. (2002). PCR-based identification of bacteria associated with endodontic infections. *J Clin Microbiol* 40:3223-3231.
- Fredricks DN, Marrazzo JM (2005). Molecular methodology in determining vaginal flora in health and disease: its time has come. *Curr Infect Dis Rep* 7:463-470.
- Fredricks DN, Relman DA (1996). Sequence-based identification of microbial pathogens: a reconsideration of Koch's postulates. *Clin Microbiol Rev* 9:18-33.
- Fredricks DN, Relman DA (1999). Application of polymerase chain reaction to the diagnosis of infectious diseases. *Clin Infect Dis* 29:475-486.
- Fredricks DN, Fiedler TL, Marrazzo JM (2005). Molecular identification of bacteria associated with bacterial vaginosis. *N Engl J Med* 353:1899-1911.
- Furrie E (2006). A molecular revolution in the study of intestinal microflora. *Gut* 55:141-143.
- Gewin V (2006). Genomics: discovery in the dirt. *Nature* 439:384-386.
- Godon JJ, Morinière J, Moletta M, Gaillac M, Bru V, Delgènes J-P (2005). Rarity associated with specific ecological niches in the bacterial world: the 'Synergistes' example. *Environ Microbiol* 7:213-224.
- Gomes BP, Lilley JD, Drucker DB (1996). Clinical significance of dental root canal microflora. *J Dent* 24:47-55.
- Gomes BP, Jacinto RC, Pinheiro ET, Sousa EL, Zaia AA, Ferraz CC, et al. (2006). Molecular analysis of *Filifactor alocis*, *Tannerella forsythia*, and *Treponema denticola* associated with primary endodontic infections and failed endodontic treatment. *J Endod* 32:937-940.
- Haapasalo M, Ranta H, Ranta K, Shah H (1986). Black-pigmented *Bacteroides* spp. in human apical periodontitis. *Infect Immun* 53:149-153.
- Handelsman J (2004). Metagenomics: application of genomics to uncultured microorganisms. *Microbiol Mol Biol Rev* 68:669-685.
- Hayashi H, Takahashi R, Nishi T, Sakamoto M, Benno Y (2005). Molecular analysis of jejunal, ileal, caecal and recto-sigmoidal human colonic microbiota using 16S rRNA gene libraries and terminal restriction fragment length polymorphism. *J Med Microbiol* 54(Pt 11):1093-1101.
- Hold GL, Pryde SE, Russel VJ, Furrie E, Flint HJ (2002). Assessment of microbial diversity in human colonic samples by 16S rDNA sequence analysis. *FEMS Microbiol Ecol* 39:33-39.
- Hughenoltz P (2002). Exploring prokaryotic diversity in the genomic era. *Genome Biol* 3:REVIEWS0003.
- Huston MA (1994). Biological diversity. Cambridge, UK: Cambridge University Press.
- Huyghe A, Francois P, Charbonnier Y, Tangomo-Bento M, Bonetti EJ, Paster BJ, et al. (2008). Novel microarray design strategy to study complex bacterial communities. *Appl Environ Microbiol* 74:1876-1885.
- Janssen PH (2006). Identifying the dominant soil bacterial taxa in libraries of 16S rRNA and 16S rRNA genes. *Appl Environ Microbiol* 72:1719-1728.
- Jung IY, Choi BK, Kum KY, Roh BD, Lee SJ, Lee CY, et al. (2000). Molecular epidemiology and association of putative pathogens in root canal infection. *J Endod* 26:599-604.
- Takehashi S, Stanley HR, Fitzgerald RJ (1965). The effects of surgical exposures of dental pulps in germ-free and conventional laboratory rats. *Oral Surg Oral Med Oral Pathol* 20:340-349.
- Kantz WE, Henry CA (1974). Isolation and classification of anaerobic bacteria from intact pulp chambers of non-vital teeth in man. *Arch Oral Biol* 19:91-96.
- Kazor CE, Mitchell PM, Lee AM, Stokes LN, Loesche WJ, Dewhirst FE, et al. (2003). Diversity of bacterial populations on the tongue dorsa of patients with halitosis and healthy patients. *J Clin Microbiol* 41:558-563.
- Keijser BJ, Zaura E, Huse SM, van der Vossen JM, Schuren FH, Montijn RC, et al. (2008). Pyrosequencing analysis of the oral microflora of healthy adults. *J Dent Res* 87:1016-1020.
- Kellam P, Weiss RA (2001). How technology drives pathogen discovery. *Microbiol Today* 29:16-18.
- Khemaleelakul S, Baumgartner JC, Pruksakorn S (2002). Identification of bacteria in acute endodontic infections and their antimicrobial susceptibility. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 94:746-755.
- Kumar PS, Griffen AL, Moeschberger ML, Leys EJ (2005). Identification of candidate periodontal pathogens and beneficial species by quantitative 16S clonal analysis. *J Clin Microbiol* 43:3944-3955.
- Lawson AJ (2004). Discovering new pathogens: culture-resistant bacteria. *Methods Mol Biol* 266:305-322.
- Le Goff A, Bunetel L, Mouton C, Bonnaure-Mallet M (1997). Evaluation of root canal bacteria and their antimicrobial susceptibility in teeth with necrotic pulp. *Oral Microbiol Immunol* 12:318-322.
- Machado de Oliveira JC, Siqueira JF Jr, Alves GB, Hirata R Jr, Andrade AF (2000). Detection of *Porphyromonas endodontalis* in infected root canals by 16S rRNA gene-directed polymerase chain reaction. *J Endod* 26:729-732.
- Molander A, Reit C, Dahlén G, Kvist T (1998). Microbiological status of root-filled teeth with apical periodontitis. *Int Endod J* 31:1-7.
- Möller AJ, Fabricius L, Dahlén G, Öhman AE, Heyden G (1981). Influence on periapical tissues of indigenous oral bacteria and necrotic pulp tissue in monkeys. *Scand J Dent Res* 89:475-484.
- Munson MA, Pitt-Ford T, Chong B, Weightman A, Wade WG (2002). Molecular and cultural analysis of the microflora associated with endodontic infections. *J Dent Res* 81:761-766; errata in *J Dent Res* 82:69, 2003, and 82:247, 2003.
- Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, et al. (2001). Bacterial diversity in human subgingival plaque. *J Bacteriol* 183:3770-3783.
- Paster BJ, Falkler WA Jr, Enwonwu CO, Idigbe EO, Savage KO, Levanos VA, et al. (2002). Prevalent bacterial species and novel phylotypes in advanced noma lesions. *J Clin Microbiol* 40:2187-2191.
- Paster BJ, Olsen I, Aas JA, Dewhirst FE (2006). The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontol* 2000 42:80-87.
- Peciuleniene V, Reynaud AH, Balciuniene I, Haapasalo M (2001). Isolation of yeasts and enteric bacteria in root-filled teeth with chronic apical periodontitis. *Int Endod J* 34:429-434.
- Pei Z, Bini EJ, Yang L, Zhou M, Francois F, Blaser MJ (2004). Bacterial biota in the human distal esophagus. *Proc Natl Acad Sci USA* 101:4250-4255.
- Pinheiro ET, Gomes BP, Ferraz CC, Sousa EL, Teixeira FB, Souza-Filho FJ (2003). Microorganisms from canals of root-filled teeth with periapical lesions. *Int Endod J* 36:1-11.
- Rajilić-Stojanović M, Smidt H, de Vos WM (2007). Diversity of the human gastrointestinal tract microbiota revisited. *Environ Microbiol* 9:2125-2136.
- Rappe MS, Giovannoni SJ (2003). The uncultured microbial majority. *Annu Rev Microbiol* 57:369-394.

- Relman DA (1999). The search for unrecognized pathogens. *Science* 284:1308-1310.
- Relman DA (2002). New technologies, human-microbe interactions, and the search for previously unrecognized pathogens. *J Infect Dis* 186(Suppl 2):254-258.
- Rôças IN, Siqueira JF Jr (2005a). Species-directed 16S rRNA gene nested PCR detection of *Olsenella* species in association with endodontic diseases. *Lett Appl Microbiol* 41:12-16.
- Rôças IN, Siqueira JF Jr (2005b). Detection of novel oral species and phylogenies in symptomatic endodontic infections including abscesses. *FEMS Microbiol Lett* 250:279-285.
- Rôças IN, Siqueira JF Jr (2008). Root canal microbiota of teeth with chronic apical periodontitis. *J Clin Microbiol* 46:3599-3606.
- Rôças IN, Siqueira JF Jr, Andrade AF, Uzeda M (2003). Oral treponemes in primary root canal infections as detected by nested PCR. *Int Endod J* 36:20-26.
- Rôças IN, Siqueira JF Jr, Santos KR (2004a). Association of *Enterococcus faecalis* with different forms of periradicular diseases. *J Endod* 30:315-320.
- Rôças IN, Jung IY, Lee CY, Siqueira JF Jr (2004b). Polymerase chain reaction identification of microorganisms in previously root-filled teeth in a South Korean population. *J Endod* 30:504-508.
- Rôças IN, Siqueira JF Jr, Aboim MC, Rosado AS (2004c). Denaturing gradient gel electrophoresis analysis of bacterial communities associated with failed endodontic treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 98:741-749.
- Rolph HJ, Lennon A, Riggio MP, Saunders WP, MacKenzie D, Coldero L, et al. (2001). Molecular identification of microorganisms from endodontic infections. *J Clin Microbiol* 39:3282-3289.
- Rupf S, Kannengiesser S, Merte K, Pfister W, Sigusch B, Eschrich K (2000). Comparison of profiles of key periodontal pathogens in periodontium and endodontium. *Endod Dent Traumatol* 16:269-275.
- Sabeti M, Simon JH, Slots J (2003). Cytomegalovirus and Epstein-Barr virus are associated with symptomatic periapical pathosis. *Oral Microbiol Immunol* 18:327-328.
- Saito D, Leonardo RdeT, Rodrigues JLM, Tsai SM, Hofling JF, Gonçalves RB (2006). Identification of bacteria in endodontic infections by sequence analysis of 16S rDNA clone libraries. *J Med Microbiol* 55(Pt 1):101-107.
- Sakamoto M, Rôças IN, Siqueira JF Jr, Benno Y (2006). Molecular analysis of bacteria in asymptomatic and symptomatic endodontic infections. *Oral Microbiol Immunol* 21:112-122.
- Sakamoto M, Siqueira JF Jr, Rôças IN, Benno Y (2007). Bacterial reduction and persistence after endodontic treatment procedures. *Oral Microbiol Immunol* 22:19-23.
- Sakamoto M, Siqueira JF Jr, Rôças IN, Benno Y (2008). Molecular analysis of the root canal microbiota associated with endodontic treatment failures. *Oral Microbiol Immunol* 23:275-281.
- Sedgley C, Nagel A, Dahlén G, Reit C, Molander A (2006). Real-time quantitative polymerase chain reaction and culture analyses of *Enterococcus faecalis* in root canals. *J Endod* 32:173-177.
- Seol JH, Cho BH, Chung CP, Bae KS (2006). Multiplex polymerase chain reaction detection of black-pigmented bacteria in infections of endodontic origin. *J Endod* 32:110-114.
- Siqueira JF Jr (2002). Endodontic infections: concepts, paradigms, and perspectives. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 94:281-293.
- Siqueira JF Jr (2008). Microbiology of apical periodontitis. In: Essential endodontology. Ørstavik D, Pitt Ford T, editors. Oxford, UK: Blackwell Munksgaard Ltd, pp. 135-196.
- Siqueira JF Jr, Rôças IN (2003a). Detection of *Filifactor alocis* in endodontic infections associated with different forms of periradicular diseases. *Oral Microbiol Immunol* 18:263-265.
- Siqueira JF Jr, Rôças IN (2003b). *Campylobacter gracilis* and *Campylobacter rectus* in primary endodontic infections. *Int Endod J* 36:174-180.
- Siqueira JF Jr, Rôças IN (2003c). *Bacteroides forsythus* in primary endodontic infections as detected by nested PCR. *J Endod* 29:390-393.
- Siqueira JF Jr, Rôças IN (2003d). *Pseudoramibacter alactolyticus* in primary endodontic infections. *J Endod* 29:735-738.
- Siqueira JF Jr, Rôças IN (2003e). Polymerase chain reaction detection of *Propionibacterium propionicus* and *Actinomyces radicidentis* in primary and persistent endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 96:215-222.
- Siqueira JF Jr, Rôças IN (2004a). Polymerase chain reaction-based analysis of microorganisms associated with failed endodontic treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 97:85-94.
- Siqueira JF Jr, Rôças IN (2004b). *Treponema* species associated with abscesses of endodontic origin. *Oral Microbiol Immunol* 19:336-339.
- Siqueira JF Jr, Rôças IN (2005a). Exploiting molecular methods to explore endodontic infections: Part 1—Current molecular technologies for microbiological diagnosis. *J Endod* 31:411-423.
- Siqueira JF Jr, Rôças IN (2005b). Exploiting molecular methods to explore endodontic infections: Part 2—Redefining the endodontic microbiota. *J Endod* 31:488-498.
- Siqueira JF Jr, Rôças IN (2005c). Uncultivated phylotypes and newly named species associated with primary and persistent endodontic infections. *J Clin Microbiol* 43:3314-3319.
- Siqueira JF Jr, Rôças IN (2006). *Catonella morbi* and *Granulicatella adiacens*: new species in endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 102:259-264.
- Siqueira JF Jr, Rôças IN (2009a). The microbiota of acute apical abscesses. *J Dent Res* 88:61-65.
- Siqueira JF Jr, Rôças IN (2009b). Community as the unit of pathogenicity: an emerging concept as to the microbial pathogenesis of apical periodontitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 107:870-878.
- Siqueira JF Jr, Rôças IN, Favieri A, Santos KR (2000a). Detection of *Treponema denticola* in endodontic infections by 16S rRNA gene-directed polymerase chain reaction. *Oral Microbiol Immunol* 15:335-337.
- Siqueira JF Jr, Rôças IN, Souto R, de Uzeda M, Colombo AP (2000b). Checkerboard DNA-DNA hybridization analysis of endodontic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 89:744-748.
- Siqueira JF Jr, Rôças IN, Oliveira JC, Santos KR (2001a). Molecular detection of black-pigmented bacteria in infections of endodontic origin. *J Endod* 27:563-566.
- Siqueira JF Jr, Rôças IN, Souto R, Uzeda M, Colombo AP (2001b). Microbiological evaluation of acute periradicular abscesses by DNA-DNA hybridization. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 92:451-457.
- Siqueira JF Jr, Rôças IN, Souto R, de Uzeda M, Colombo AP (2002). *Actinomyces* species, streptococci, and *Enterococcus faecalis* in primary root canal infections. *J Endod* 28:168-172.
- Siqueira JF Jr, Rôças IN, Andrade AF, de Uzeda M (2003). *Peptostreptococcus micros* in primary endodontic infections as detected by 16S rDNA-based polymerase chain reaction. *J Endod* 29:111-113.
- Siqueira JF Jr, Rôças IN, Alves FR, Santos KR (2004a). Selected endodontic pathogens in the apical third of infected root canals: a molecular investigation. *J Endod* 30:638-643.
- Siqueira JF Jr, Rôças IN, Rosado AS (2004b). Investigation of bacterial communities associated with asymptomatic and symptomatic endodontic infections by denaturing gradient gel electrophoresis fingerprinting approach. *Oral Microbiol Immunol* 19:363-370.
- Siqueira JF Jr, Rôças IN, Cunha CD, Rosado AS (2005). Novel bacterial phylotypes in endodontic infections. *J Dent Res* 84:565-569.
- Siqueira JF Jr, Paiva SS, Rôças IN (2007a). Reduction in the cultivable bacterial populations in infected root canals by a chlorhexidine-based antimicrobial protocol. *J Endod* 33:541-547.
- Siqueira JF Jr, Magalhães KM, Rôças IN (2007b). Bacterial reduction in infected root canals treated with 2.5% NaOCl as an irrigant and calcium hydroxide/camphorated paramonochlorophenol paste as an intracanal dressing. *J Endod* 33:667-672.
- Siqueira JF Jr, Rôças IN, Paiva SSM, Magalhães KM, Guimarães-Pinto T (2007c). Cultivable bacteria in infected root canals as identified by 16S rRNA gene sequencing. *Oral Microbiol Immunol* 22:266-271.
- Siqueira JF Jr, Rôças IN, Paiva SS, Guimarães-Pinto T, Magalhães KM, Lima KC (2007d). Bacteriologic investigation of the effects of sodium hypochlorite and chlorhexidine during the endodontic treatment of teeth with apical periodontitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 104:122-130.

- Sjögren U, Figdor D, Persson S, Sundqvist G (1997). Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *Int Endod J* 30:297-306.
- Sogin ML, Morrison HG, Huber JA, Welch DM, Huse SM, Neal PR, et al. (2006). Microbial diversity in the deep sea and the underexplored "rare biosphere". *Proc Natl Acad Sci USA* 103:12115-12120.
- Spiegelman D, Whissell G, Greer CW (2005). A survey of the methods for the characterization of microbial consortia and communities. *Can J Microbiol* 51:355-386.
- Suau A, Bonnet R, Sutren M, Godon JJ, Gibson GR, Collins MD, et al. (1999). Direct analysis of genes encoding 16S rRNA from complex communities reveals many novel molecular species within the human gut. *Appl Environ Microbiol* 65:4799-4807.
- Sunde PT, Tronstad L, Eribe ER, Lind PO, Olsen I (2000). Assessment of periradicular microbiota by DNA-DNA hybridization. *Endod Dent Traumatol* 16:191-196.
- Sundqvist G (1976). Bacteriological studies of necrotic dental pulps [Odontological Dissertation no. 7]. Umeå, Sweden: University of Umeå, Odontological Dissertations, pp. 1-93.
- Sundqvist G (1992). Associations between microbial species in dental root canal infections. *Oral Microbiol Immunol* 7:257-262.
- Sundqvist G, Johansson E, Sjögren U (1989). Prevalence of black-pigmented bacteroides species in root canal infections. *J Endod* 15:13-19.
- Sundqvist G, Figdor D, Persson S, Sjogren U (1998). Microbiologic analysis of teeth with failed endodontic treatment and the outcome of conservative re-treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 85:86-93.
- Tronstad L, Sunde PT (2003). The evolving new understanding of endodontic infections. *Endod Topics* 6:57-77.
- Tronstad L, Barnett F, Riso K, Slots J (1987). Extraradicular endodontic infections. *Endod Dent Traumatol* 3:86-90.
- van Winkelhoff AJ, Carlee AW, de Graaff J (1985). *Bacteroides endodontalis* and other black-pigmented *Bacteroides* species in odontogenic abscesses. *Infect Immun* 49:494-498.
- Vianna ME, Conrads G, Gomes BPFA, Horz HP (2006a). Identification and quantification of archaea involved in primary endodontic infections. *J Clin Microbiol* 44:1274-1282.
- Vianna ME, Horz HP, Gomes BP, Conrads G (2006b). *In vivo* evaluation of microbial reduction after chemo-mechanical preparation of human root canals containing necrotic pulp tissue. *Int Endod J* 39:484-492.
- Vianna ME, Conrads G, Gomes BPFA, Horz HP (2007). Quantification and characterization of *Synergistes* in endodontic infections. *Oral Microbiol Immunol* 22:260-265.
- Vickerman MM, Brossard KA, Funk DB, Jesionowski AM, Gill SR (2007). Phylogenetic analysis of bacterial and archaeal species in symptomatic and asymptomatic endodontic infections. *J Med Microbiol* 56:110-118.
- von Wintzingerode F, Gobel UB, Stackebrandt E (1997). Determination of microbial diversity in environmental samples: pitfalls of PCR-based rRNA analysis. *FEMS Microbiol Rev* 21:213-229.
- Weiger R, Manncke B, Werner H, Lost C (1995). Microbial flora of sinus tracts and root canals of non-vital teeth. *Endod Dent Traumatol* 11:15-19.
- Wittgow WC Jr, Sabiston CB Jr (1975). Microorganisms from pulpal chambers of intact teeth with necrotic pulps. *J Endod* 1:168-171.
- Zoletti GO, Siqueira JF Jr, Santos KR (2006). Identification of *Enterococcus faecalis* in root-filled teeth with or without periradicular lesions by culture-dependent and -independent approaches. *J Endod* 32:722-726.