

The evaluation of removal of calcium hydroxide paste from an artificial standardized groove in the apical root canal using different irrigation methodologies

L. W. M. van der Sluis, M. K. Wu & P. R. Wesselink

Department of Cariology Endodontology Pedodontology, Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, The Netherlands

Abstract

van der Sluis LWM, Wu MK, Wesselink PR. The evaluation of removal of calcium hydroxide paste from an artificial standardized groove in the apical root canal using different irrigation methodologies. *International Endodontic Journal*, **40**, 52–57, 2007.

Aim To evaluate the capacity to remove a calcium hydroxide ($\text{Ca}(\text{OH})_2$) paste from the root canal and to evaluate the efficacy of $\text{Ca}(\text{OH})_2$ removal during passive ultrasonic irrigation using either sodium hypochlorite (NaOCl) or water as an irrigant.

Methodology Sixteen mandibular premolars were used. Each root was prepared to the apical foramen using GT instruments of size 30, 0.06 taper. Each root was split longitudinally. In one half of the root, a groove was cut in the canal wall 2–6 mm from the apex which was then filled with a $\text{Ca}(\text{OH})_2$ paste. Subsequently the roots were reassembled. In group 1 ($n = 16$), the teeth were ultrasonically irrigated using 50 mL 2.0% NaOCl as the irrigant. Group 2 ($n = 16$) was treated in the same manner as group 1, but using

50 mL water in place of the NaOCl . In group 3 ($n = 16$), the teeth were irrigated by syringe injection of 50 mL 2.0% NaOCl . The quantity of remaining $\text{Ca}(\text{OH})_2$ in the groove was scored and the data analysed with Kruskal–Wallis and Mann–Whitney tests.

Results The difference in remaining $\text{Ca}(\text{OH})_2$ between all groups was statistically significant ($P < 0.001$). Group 1 had significantly lower scores than group 2 ($P < 0.001$) and group 3 ($P = 0.002$), but there was no significant difference between groups 2 and 3 ($P = 0.765$).

Conclusions Passive ultrasonic irrigation with 2% NaOCl was more effective in removing $\text{Ca}(\text{OH})_2$ paste from artificial root canal grooves than syringe delivery of 2% NaOCl or water as irrigant.

Keywords: calcium hydroxide, passive ultrasonic irrigation, sodium hypochlorite, syringe delivery, water.

Received 10 March 2006; accepted 19 June 2006

Introduction

Calcium hydroxide ($\text{Ca}(\text{OH})_2$) in a nonsetting paste form is used widely as an intracanal medicament during root canal treatment because of its antimicrobial properties (Siqueira & Lopes 1999, Law & Messer

2004). The most frequently described method for removal of $\text{Ca}(\text{OH})_2$ from the root canal is instrumentation of the root canal with the master apical file in combination with copious irrigation of sodium hypochlorite (NaOCl) and EDTA (Lambrianidis *et al.* 2006). EDTA has the ability to neutralize $\text{Ca}(\text{OH})_2$ residues, which could prevent a chemical reaction with the sealer, or chelate $\text{Ca}(\text{OH})_2$ residues, which could make them easier to remove by irrigation (Margelos *et al.* 1997). However, it has been reported that removal of $\text{Ca}(\text{OH})_2$ from the apical root canal wall, when this

Correspondence: L. W. M. van der Sluis, Department of Cariology Endodontology Pedodontology, ACTA, Louwesweg 1, 1066 EA Amsterdam, The Netherlands (Tel.: +3120 518 8651; fax: +3120 669 2881; e-mail: lvd.sluis@acta.nl).

method is used, is difficult (Margelos *et al.* 1997, Lambrianidis *et al.* 1999, 2006). This can be explained because instrumentation and irrigation alone cannot completely clean the entire root canal wall (Wu *et al.* 2003). When Ca(OH)₂ is removed from the main canal with a file, remnants will remain in canal extensions or irregularities. From these canal extensions or irregularities it is only possible to remove the Ca(OH)₂ by irrigation. Passive ultrasonic irrigation (PUI) is more effective in dentine debris removal from the root canal wall than syringe delivery of the irrigant (Goodman *et al.* 1985, Lee *et al.* 2004b). However, whether PUI can remove Ca(OH)₂ from the root canal wall is not known.

Passive ultrasonic irrigation is based on the transmission of energy from an ultrasonically oscillating instrument to the irrigant in the root canal (van der Sluis *et al.* 2005a). In the irrigant solution, acoustic streaming and/or cavitation will occur that is more powerful when the root canal wall is not touched deliberately with the oscillating instrument (Ahmad *et al.* 1988, Roy *et al.* 1994). Therefore, PUI is undertaken after the root canal is prepared up to the master apical file and without instrumenting the root canal wall during the ultrasonic irrigation.

Calcium hydroxide remnants left in the root canal can result in a thicker nonhomogenous appearance of root canal sealers (Margelos *et al.* 1997, Kim & Kim 2002, Hosoya *et al.* 2004). The sealer thickness could have an effect on the sealing ability of root canal fillings (Kontakiotis *et al.* 1997). The Ca(OH)₂ remnants could also result in a chemical reaction with the sealer resulting in a reduction of flow or working time (Margelos *et al.* 1997, Hosoya *et al.* 2004). Cracks were visible in zinc-oxide eugenol sealer after the use of Ca(OH)₂ (Kim & Kim 2002) which can be explained by the faster setting of the sealer under the influence of Ca(OH)₂ (Margelos *et al.* 1997). Calcium hydroxide remnants could also prevent sealer from penetrating the dentinal tubules resulting in a potential reduction in sealer adaption (Çalt & Serper 1999). The dimensional instability of Ca(OH)₂ and its potential to dissolve in water and dissociate into hydroxide and calcium ions (Cohen & Burns 2002) could influence the leakage of root fillings on the long term. In the past, leakage studies have been undertaken to evaluate the influence of the above-mentioned phenomena on the leakage of a root canal filling *ex vivo*. However, most of these were passive dye leakage studies which are not reliable because Ca(OH)₂ discolours methylene blue (Wu *et al.* 1998), which is often used as a colouring agent for dye

leakage. Furthermore, dye leakage is a passive leakage test which can hinder the penetration of dye when air bubbles are present between the root filling and the root canal wall (Wu *et al.* 1994). Because the studies were performed directly after filling the root canals, the effect on the long term could not be evaluated. Therefore, a negative effect of Ca(OH)₂ remnants on leakage of subsequently placed root fillings on the short and long term has never been demonstrated.

In a study by Lee *et al.* (2004a), a standardized groove was cut in the apical part of the root canal to simulate an oval extension of the apical root canal. The dimensions of the groove were determined using the data of the anatomy of the apical root canal as described by Wu *et al.* (2000). Using this methodology the removal of calcium hydroxide paste packed in an apical extension, which cannot be removed by endodontic files, but can only be removed by irrigation, could be evaluated while the situation before irrigation is controlled.

The purpose of the study was to evaluate the capacity to remove Ca(OH)₂ from an artificial standardized groove in the apical root canal and to evaluate the efficacy of Ca(OH)₂ removal during PUI between NaOCl and water.

Materials and methods

Sixteen mandibular single-rooted premolars were selected after bucco-lingual and mesio-distal radiographs indicated that their internal diameters at three points (2, 4 and 6 mm from the apex) were smaller than the corresponding diameters of a size 30, 0.06 taper GT instrument (Dentsply Maillefer, Ballaigues, Switzerland). The crowns of the teeth were removed 12 mm from the apex to standardize the length of the roots and eliminate the variation in dimension of the pulp chamber. The root canals were shaped with GT root canal instruments, to a size 30, 0.06 taper GT instrument as the master apical file. Each canal was prepared to the apical foramen which was determined by inserting a size 10 K-file into the canal until the tip of the file was just visible. Between each instrument, the canals were irrigated with 2 mL of a freshly prepared 2% solution of NaOCl, using a syringe and a 27-gauge needle that was placed 1 mm short of the working length, resulting in a total volume of 30 mL. The NaOCl solution was prepared by diluting a 10% NaOCl solution (Merck, Darmstadt, Germany). Its pH was adjusted to 10.8 with 1 N HCl. The concentration of the NaOCl solution was measured iodometrically (Moorer & Wesselink 1982).

After root canal preparation each root was split longitudinally through the canal, forming two halves. A standard groove of 4 mm in length, 0.2 mm in width and 0.5 mm in depth was cut in one canal wall 2–6 mm from the apex, to simulate uninstrumented canal extensions in the apical half (Lee et al. 2004a) (Fig. 1). Each groove was filled with calcium hydroxide paste (Ultracal[®] XS; Ultradent products, Jordan, UT, USA) using paper points 30–40, to simulate a situation when calcium hydroxide paste remains in natural canal extensions after instrumentation. Care was taken to consistently apply the calcium hydroxide paste in the groove. The teeth were reassembled with wires and stored for 1 week at 37 °C in 100% relative humidity to simulate the clinical situation when Ca(OH)₂ is used as an intermediate root filling between two treatment visits. After 1 week of storage, images of each half of the canal with a groove were taken using a Photomakroskop M 400 microscope with a digital camera (Wild, Heerbrugg, Switzerland) at ×40 magnification to detect the amount of Ca(OH)₂ just before irrigation; the photos were then scanned as tagged-image file format images. The two root halves were reassembled by means of wires and sticky wax, and the apical foramen was sealed with wax. Three different types of irrigation were tested with the 16 teeth; a pilot study had demonstrated that the smooth wire used during ultrasonic irrigation did not damage the root dentine or alter the form of the groove. In group 1 (*n* = 16), the root canals were ultrasonically irrigated for 3 min with a continuous flow of 50 mL 2.0% NaOCl. Group 2 (*n* = 16) was the same as group 1 with the exception that water was used




Figure 1 Schematic representation of specimen preparation. In one half of the instrumented root canal, a groove was cut 2–6 mm from the apex.

as the irrigant in place of 2% NaOCl. In group 3 (*n* = 16), the canals were irrigated with 50 mL of 2.0% NaOCl using a syringe with a 27-gauge Terumo needle (Terumo Europe N.V., Brussels, Belgium) which was inserted just short of the apical foramen. Ultrasonic irrigation was performed with a piezoelectronic unit (PMax; Satelec, Merignac, France). After switching on the ultrasonic device, an activated 21-mm-long stainless steel smooth wire with a diameter of 0.15 mm and a 0.02 taper (van der Sluis et al. 2005a) was placed in the canal just short of the apical foramen, the oscillation of the wire and irrigation began almost at the same time. The oscillation of the wire was directed towards the groove and the intensity was set on speed 'blue 4'. According to the manufacturer, the frequency employed under these conditions was approximately 30 kHz, and the displacement-amplitude varied between 20 and 30 μm. Between the different irrigation procedures the root halves containing the groove were cleaned with brushes and air under high pressure. The grooves were examined under a microscope to ensure that all Ca(OH)₂ remnants were removed from the grooves. The root halves were separated after each irrigation procedure to evaluate the removal of Ca(OH)₂. After irrigation, images of each half of the canal with a groove were taken using a Photomakroskop M 400 microscope with a digital camera (Wild) at ×40 magnification; the photos were then scanned as tagged-image file format images.

The quantity of Ca(OH)₂ in the groove before and after irrigation was scored double-blind and independently by three calibrated dentists using the following scores: score 0, the groove is empty; score 1, less than half of the groove is filled with Ca(OH)₂; score 2, more than half of the groove is filled with Ca(OH)₂ and score 3, the groove is filled completely with Ca(OH)₂. With the scores before and after irrigation, the percentage of score reduction was calculated as follows:

$$\text{Percentage of score reduction} = \frac{\text{score before irrigation} - \text{score after irrigation}}{\text{score before irrigation}} \times 100\%.$$

The differences in Ca(OH)₂ scores between the different groups were analysed by means of the Kruskal–Wallis and Mann–Whitney tests. The level of significance was set at $\alpha = 0.05$.

Results

The results of the study are shown in Table 1. The Ca(OH)₂ score was reduced by 63.3% in group 1, 6.7%

Table 1 Score of calcium hydroxide before and after irrigation and the percentage of score reduction

Groups	Before	After		Percentage of score reduction	
		Mean	SD	Mean	SD
1	3.00	1.1	1.1	63.3	36.7
2	3.00	2.8	0.5	6.7	16.7
3	3.00	2.5	1.0	16.7	33

in group 2 and 16.7% in group 3. The difference between all groups was statistically significant (Kruskal–Wallis test, $P < 0.001$). Group 1 differed significantly from group 2 ($P < 0.001$) and group 3 ($P = 0.002$), but there was no significant difference between groups 2 and 3 ($P = 0.765$).

Discussion

The results indicate that PUI with NaOCl as irrigant was more effective in removal of $\text{Ca}(\text{OH})_2$ paste from an artificial standardized groove in the apical root canal than PUI with water or irrigation by means of syringe delivery of NaOCl.

During PUI, acoustic microstreaming and cavitation can occur which cause a streaming pattern within the root canal from the apical to the coronal (Ahmad *et al.* 1987, Roy *et al.* 1994). Because of this microstreaming, more dentine debris can be removed from the root canal compared with syringe delivery of the irrigant (Lee *et al.* 2004a), even from remote places in the root canal (Goodman *et al.* 1985). Probably the same mechanisms are responsible for the more effective removal of $\text{Ca}(\text{OH})_2$ during PUI in comparison with syringe delivery of the irrigant.

Sodium hypochlorite as irrigant is more effective in removing dentine debris from the root canal during PUI than water (Huque *et al.* 1998, van der Sluis *et al.* 2006). As 30% of dentine debris is an organic material, the excellent organic tissue dissolution properties of NaOCl were suggested as an explanation (van der Sluis *et al.* 2006). To examine if the organic tissue dissolution properties are really of importance for the effectiveness of PUI with NaOCl as irrigant, the removal of a nonorganic substance from a standardized groove in a root canal during PUI could give an answer. The capacity to remove a nonorganic substance from a standardized groove in a root canal during PUI can evaluate the importance of the tissue-dissolving effect of NaOCl during PUI. Sodium hypochlorite as an irrigant during PUI is also more effective

in removing $\text{Ca}(\text{OH})_2$ from an artificial groove in the apical root canal than water, as it is during the removal of dentine debris. The results of PUI with water were comparable with the results of irrigation by means of syringe delivery of NaOCl indicating that the extra capacity to remove matter from the root canal of PUI with NaOCl as irrigant does not occur when water is used. Because $\text{Ca}(\text{OH})_2$ is an inorganic substance the tissue dissolving capacity of NaOCl cannot play a role in the process. There are several explanations for differences between water and NaOCl as irrigant during PUI. First the physical properties of NaOCl are different from water, NaOCl is a salt water suspension, normal water not. Bubbles formed in salt water tend to be more numerous, particularly regarding the smallest bubbles and are less prone to coalesce than bubbles in fresh water (Leighton 1994). Because the smallest bubbles are more numerous, the acoustic microstreaming will be different and could perhaps be more powerful. Another explanation is that gas will dissolve in the bubble during cavitation and the oscillations of the bubble depend on the concentration of the gas dissolved in the liquid, the temperature of the liquid and small amounts of surface active impurities (Brenner *et al.* 2002). During PUI with NaOCl as irrigant, chlorine gas will be present in the irrigant which can dissolve in the bubble. The chlorine gas in the irrigant will also have an influence on the oscillation of the bubbles. This will influence the acoustic microstreaming. Whether the temperature and surface active impurities during PUI are different with NaOCl or water as an irrigant is not known. All these items remain for further research.

In this study, the most successful method of $\text{Ca}(\text{OH})_2$ removal was PUI with NaOCl as irrigant; an average of 63.3% of $\text{Ca}(\text{OH})_2$ removal was recorded. In earlier studies, higher percentages of dentine debris removal and more samples with a completely empty groove were recorded (Lee *et al.* 2004b, van der Sluis *et al.* 2005b, 2006) (Table 2). However, root canals of different sizes and tapers were used varying from apical sizes 50, 30 and 20 and tapers of 0.10–0.06. It is more difficult to remove dentine debris from the root canal when the size and taper of the root canal are smaller (Lee *et al.* 2004a, van der Sluis *et al.* 2005b). However, the average removal of $\text{Ca}(\text{OH})_2$ from a size 30 taper 0.06 was even lower than the scores of dentine debris removal from a smaller root canal of size 20 and taper 0.06. This could indicate that it is more difficult to remove $\text{Ca}(\text{OH})_2$ from the root canal wall than dentine debris. The remaining 36.7% $\text{Ca}(\text{OH})_2$ residue could

Table 2 Percentage of dentine debris or Ca(OH)₂ removal from an artificial groove in the apical root canal in different studies

Study	Size	Taper	Irrigation	Reduction (%)	Score 0 (%)
Lee et al. (2004b), dentine debris	50	SB	US NaOCl	88	75
	50	SB	S NaOCl	25	0
van der Sluis et al. (2005b), dentine debris	20	0.10	US NaOCl	92.7	87
van der Sluis et al. (2006), dentine debris	20	0.10	US NaOCl	98	93
	20	0.10	US water	44	20
Ca(OH) ₂	30	0.08	US NaOCl	63.3	37.5
	30	0.08	US water	6.7	0
	30	0.08	S NaOCl	16.7	6.25

have a negative effect on the sealing abilities of sealers, prevent sealers from penetrating the dentinal tubules or dissolve with time (Margelos et al. 1997, Çalt & Serper 1999, Cohen & Burns 2002, Kim & Kim 2002, Hosoya et al. 2004).

No irrigation with EDTA was used in this study. EDTA may neutralize Ca(OH)₂ residues to prevent chemical influence on the sealer; however, the interference from the mechanical point of view is still present (Margelos et al. 1997). Çalt & Serper (1999) reported complete removal of Ca(OH)₂ from the root canal after irrigation with EDTA and NaOCl in comparison with NaOCl alone. It is likely that EDTA may chelate residual Ca(OH)₂ which is then more easily removed by irrigation with NaOCl (Margelos et al. 1997). However, other studies using the same irrigation regime (EDTA and NaOCl) could not confirm these results and still found extensive remnants of Ca(OH)₂ (Tatsuta et al. 1999, Lambrianidis et al. 2006). There is no evidence that EDTA can completely dissolve Ca(OH)₂ placed superficially on the canal wall or from deeper layers of the root canal. However, it could be interesting to study if pre-treatment of Ca(OH)₂ with EDTA removes Ca(OH)₂ more easily from the root canal during PUI than no pre-treatment of EDTA.

In this study, Ultracal[®] XS was used as the calcium hydroxide paste. UltraCal[®] XS is an aqueous radiopaque paste with a pH of 12.5. The approximate Ca(OH)₂ concentration is 35%. If another type of calcium hydroxide paste would have given different results is not known, but as this is an aqueous solution of Ca(OH)₂ no influence from the vehicle is expected.

Conclusion

Passive ultrasonic irrigation with 2% NaOCl is more effective in removing Ca(OH)₂ from the root canal than syringe delivery of 2% NaOCl or water as irrigant during PUI.

References

- Ahmad M, Pitt Ford TR, Crum LA (1987) Ultrasonic debridement of root canals: acoustic streaming and its possible role. *Journal of Endodontics* **14**, 490–9.
- Ahmad M, Pitt Ford TR, Crum LA, Walton AJ (1988) Ultrasonic debridement of rootcanals: acoustic cavitation and its relevance. *Journal of Endodontics* **14**, 486–93.
- Brenner MP, Hilgenfeldt S, Lohse D (2002) Single bubble sonoluminescence. *Reviews of modern physics* **74**, 425–84.
- Çalt S, Serper A (1999) Dentinal tubule penetration of root canal sealers after root canal dressing with calcium hydroxide. *Journal of Endodontics* **25**, 431–3.
- Cohen S, Burns RC (2002) *Pathways of the Pulp*, 8th edn. St Louis, MO, USA: Mosby Inc., p. 548.
- Goodman A, Reader A, Beck M, Melfi R, Meyers W (1985) An *in vitro* comparison of the efficacy of the step-back technique versus a step-back/ultrasonic technique in human mandibular molars. *Journal of Endodontics* **11**, 249–56.
- Hosoya N, Kurayama H, Lino F, Arai T (2004) Effects of calcium hydroxide on physical and sealing properties of canal sealers. *International Endodontic Journal* **37**, 178–84.
- Huque J, Kota K, Yamaga M, Iwaku M, Hoshino E (1998) Bacterial eradication from root dentine by ultrasonic irrigation with sodium hypochlorite. *International Endodontic Journal* **31**, 242–50.
- Kim SK, Kim YO (2002) Influence of calcium hydroxide intracanal medication on apical seal. *International Endodontic Journal* **35**, 623–8.
- Kontakiotis EG, Wu MK, Wesselink PR (1997) Effect of sealer thickness on long-term sealing ability: a 2-year follow-up study. *International Endodontic Journal* **30**, 307–12.
- Lambrianidis T, Margelos J, Beltes P (1999) Removal efficacy of calcium hydroxide dressing from the root canal. *Journal of Endodontics* **25**, 85–8.
- Lambrianidis T, Kosti E, Boutsoukias C, Mazinis M (2006) Removal efficacy of various calcium hydroxide/chlorhexidine medicaments from the root canal. *International Endodontic Journal* **39**, 55–61.
- Law A, Messer H (2004) An evidence-based analysis of the antimicrobial effectiveness of intracanal medicaments. *Journal of Endodontics* **30**, 689–94.

- Lee S-J, Wu M-K, Wesselink PR (2004a) The efficacy of ultrasonic irrigation to remove artificially placed dentine debris from different-sized simulated plastic root canals. *International Endodontic Journal* **37**, 607–12.
- Lee S-J, Wu M-K, Wesselink PR (2004b) The effectiveness of syringe irrigation and ultrasonics to remove debris from simulated irregularities within prepared root canal walls. *International Endodontic Journal* **37**, 672–8.
- Leighton TG (1994) *The Acoustic Bubble*, Chapters 1 and 2. New York: Academic Press.
- Margelos J, Eliades G, Verdelis C, Palagias G (1997) Interaction of calcium hydroxide with zinc oxide–eugenol type sealers: a potential clinical problem. *Journal of Endodontics* **23**, 43–51.
- Moorer WR, Wesselink PR (1982) Factors promoting the tissue dissolving capability of sodium hypochlorite. *International Endodontic Journal* **15**, 187–96.
- Roy RA, Ahmad M, Crum LA (1994) Physical mechanisms governing the hydrodynamic response of an oscillating ultrasonic file. *International Endodontic Journal* **27**, 197–207.
- Siqueira LF, Lopes HP (1999) Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *International Endodontic Journal* **32**, 361–9.
- van der Sluis LWM, Wu MK, Wesselink PR (2005a) A comparison between a smooth wire and a K file in the removal of artificially placed dentine debris in plastic blocks. *International Endodontic Journal* **38**, 593–6.
- van der Sluis LWM, Wu MK, Wesselink PR (2005b) The efficacy of ultrasonic irrigation to remove artificially placed dentine debris from human root canals prepared using instruments of varying taper. *International Endodontic Journal* **38**, 764–8.
- van der Sluis LWM, Wu MK, Wesselink PR (2006) The influence of volume, type of irrigant and flushing method on removing artificially placed dentine debris from the apical root canal during passive ultrasonic irrigation. *International Endodontic Journal* **39**, 472–6.
- Tatsuta CT, Morgan LA, Baumgartner JC, Adey JD (1999) Effect of calcium hydroxide and four irrigation regimes on instrumented and uninstrumented canal wall topography. *Journal of Endodontics* **25**, 93–8.
- Wu MK, de Gee AJ, Wesselink PR (1994) Fluid transport and dye penetration along root canal fillings. *International Endodontic Journal* **27**, 233–8.
- Wu MK, Kontakiotis EG, Wesselink PR (1998) Decoloration of 1% methylene blue solution in contact with dental filling materials. *Journal of Dentistry* **26**, 585–9.
- Wu MK, Roris A, Barkis D, Wesselink PR (2000) Prevalence and extent of long oval canals in the apical third. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontology* **89**, 739–43.
- Wu MK, van der Sluis LWM, Wesselink PR (2003) The capability of two hand instrumentation techniques to remove the inner layer of dentine in oval canals. *International Endodontic Journal* **36**, 218–24.