

REVIEW

Limitations of previously published systematic reviews evaluating the outcome of endodontic treatment

M-K. Wu, H. Shemesh & P. R. Wesselink

Department of Endodontology, Academic Centre of Dentistry Amsterdam (ACTA), University of Amsterdam and VU University, Amsterdam, The Netherlands

Abstract

Wu M-K, Shemesh H, Wesselink PR. Limitations of previously published systematic reviews evaluating the outcome of endodontic treatment. *International Endodontic Journal*.

The aim of this work was to identify the limitations of previously published systematic reviews evaluating the outcome of root canal treatment. Traditionally, periapical radiography has been used to assess the outcome of root canal treatment with the absence of a periapical radiolucency being considered a confirmation of a healthy periapex. However, a high percentage of cases confirmed as healthy by radiographs revealed apical periodontitis on cone beam computed tomography (CBCT) and by histology. In teeth, where reduced size of the existing radiolucency was diagnosed by radiographs and considered to represent periapical healing, enlargement of the lesion was frequently confirmed by CBCT. In clinical studies, two additional factors may have further contributed to the overestimation of successful outcomes after root canal treatment:

(i) extractions and re-treatments were rarely recorded as failures; and (ii) the recall rate was often lower than 50%. The periapical index (PAI), frequently used for determination of success, was based on radiographic and histological findings in the periapical region of maxillary incisors. The validity of using PAI for all tooth positions might be questionable, as the thickness of the cortical bone and the position of the root tip in relation with the cortex vary with tooth position. In conclusion, the serious limitations of longitudinal clinical studies restrict the correct interpretation of root canal treatment outcomes. Systematic reviews reporting the success rates of root canal treatment without referring to these limitations may mislead readers. The outcomes of root canal treatment should be re-evaluated in long-term longitudinal studies using CBCT and stricter evaluation criteria.

Keywords: outcome, root canal treatment, success, systematic review.

Received 13 January 2009; accepted 28 April 2009

Introduction

Apical periodontitis may present before and after root canal treatment. Teeth with apical periodontitis may be symptomatic or asymptomatic, functional or not functional. Ørstavik & Pitt Ford (1998), Friedman (2002)

and Trope (2003) have defined clinical endodontics as the prevention and/or elimination of apical periodontitis. Accordingly, the aim of root canal treatment is to reduce root infection to a minimal level and achieve absence of post-treatment apical periodontitis (Wu *et al.* 2006).

The outcome of root canal treatment indicates the extent to which the above aims have been achieved. In an effort to provide patients with the most recent, highest quality and most predictable treatment modalities for dental care, clinicians must be well informed

Correspondence: M-K. Wu, Department of Endodontology, ACTA, Louwesweg 1, 1066 EA Amsterdam, The Netherlands (Tel.: +31 20 5188367; fax: +31 20 6692881; e-mail: m.wu@acta.nl).

regarding the outcomes of their proposed treatment. Moreover, the selection of treatment procedures, instruments and materials is often determined based on evidence of a higher success rate. Therefore, evidence-based knowledge of root canal treatment outcomes is a fundamental aspect of endodontics.

Longitudinal clinical studies, including randomized controlled trials, cohort studies and retrospective observational studies provide information about the outcomes of initial root canal treatments and surgical or nonsurgical retreatments (Paik *et al.* 2004, Ng *et al.* 2007). Eleven systematic reviews on the outcome of root canal treatment have been published in the last decade (Table 1), summarizing findings from longitudinal clinical studies published between 1922 and 2006 (Hepworth & Friedman 1997, Peterson & Gutmann 2001, Lewsey *et al.* 2001, Basmadjian-Charles *et al.* 2002, Niederman & Theodosopoulou 2003, Paik *et al.* 2004, Kojima *et al.* 2004, Sathorn *et al.* 2005, Ng *et al.* 2008a, 2008b). Review articles can influence clinicians more than individual clinical studies, because they are considered to represent high-quality (sub-standard) evidence in the absence of sufficient gold standard level data from randomized controlled trials (Ng *et al.* 2007).

However, previously published systematic reviews on the outcome of root canal treatment may have neglected a number of crucial factors affecting the assessment of overall treatment success (Wu *et al.* 2006). One such factor is that periapical radiography was used for all individual clinical studies; this technique has limited value in diagnosing a healthy

periapex (Bender & Seltzer 1961, Bender 1982, van der Stelt 1985, Stabholz *et al.* 1994, Huumonen & Ørstavik 2002, Ricucci & Bergenholtz 2003).

The purpose of this work was to highlight the limitations of previously published systematic reviews evaluating the outcomes of root canal treatment.

Periapical radiography

In 57% of clinical studies selected in a review article by Ng *et al.* (2007), both clinical and radiographic findings were used to determine the treatment outcome. As post-treatment apical periodontitis is often asymptomatic, the outcome was determined by radiographic examination alone in the remaining 43% of selected studies. Either strict (complete resolution of existing periapical radiolucency at recall) or loose (reduction in size of existing periapical radiolucency at recall) radiographic criteria were used in these studies.

Although, periapical radiographs have been used to diagnose post-treatment apical periodontitis in all studies since 1922 (Ng *et al.* 2007), the limitations of periapical radiography in diagnosing apical periodontitis were not discussed in any of the systematic reviews listed in Table 1.

The image on radiographs corresponds to a two-dimensional aspect of a three-dimensional structure (Huumonen & Ørstavik 2002). Periapical lesions confined within the cancellous bone may not be detected by periapical radiography. Whilst lesions of a certain size can be detected in regions covered by thin cortex, lesions of the same size cannot be detected in regions

Author (year)	Country	Treatment	Estimated success (%)
Hepworth & Friedman (1997)	Canada	Retreatment	66
		Apical surgery	59
		Apical surgery with simultaneous retreatment	81
Peterson & Gutmann (2001)	USA	Resurgery	36
Lewsey <i>et al.</i> (2001)	UK	Root canal treatment	78
Basmadjian-Charles <i>et al.</i> (2002)	France	Root canal treatment	78
Niederman & Theodosopoulou (2003)	USA	Retrograde filling	77
Paik <i>et al.</i> (2004)	USA	Retreatment	70
Kojima <i>et al.</i> (2004)	Japan	Vital pulp	83
		Nonvital	79
Sathorn <i>et al.</i> (2005)	Australia	Single-visit	77
		Multiple-visit	71
Ng <i>et al.</i> (2007)	UK	Root canal treatment	75
Ng <i>et al.</i> (2008a)	UK	Root canal treatment	–
Ng <i>et al.</i> (2008b)	UK	Retreatment	77

Table 1 Estimated success rate of endodontic treatments reported in previously published systematic reviews (1997–2008)

covered by thicker cortex, which has been demonstrated in both *in vivo* and *ex vivo* studies (Bender & Seltzer 1961, Bender 1982, van der Stelt 1985, Stabholz *et al.* 1994, Huuonen & Ørstavik 2002, Ricucci & Bergenholz 2003).

Computed tomography (CT) has been widely used in medicine since the 1970s (Brenner & Hall 2007) and first appeared in endodontic research in 1990 (Tachibana & Matsumoto 1990). Cone-beam technology has existed since the 1980s. Cone beam computed tomography (CBCT) can detect periapical lesions in many cases where no periapical radiolucency can be seen on radiographs (Vandenbergh *et al.* 2008).

Lofthag-Hansen *et al.* (2007) diagnosed periapical lesions in endodontically treated human molar teeth using both periapical radiography and CBCT. Forty-six molars were inspected and 53 lesions were detected by both techniques. In addition, 33 lesions were detected by CBCT only. Jorge *et al.* (2008) infected 76 tooth roots in dogs and detected no periapical lesions with radiographs at day 14 after pulp exposure, whilst 47% of roots showed lesions at day 21. By contrast, CBCT evaluation detected apical periodontitis in 33% of roots at day 14 and 83% at day 21. Cross-sectional studies were considered by some to provide reliable information on the long-term success rate of root canal treatment at a population level (Pettersson *et al.* 1991). Estrela *et al.* (2008a) in a cross-sectional study demonstrated post-treatment apical periodontitis in 35% of teeth using periapical radiography and in 63% of teeth using CBCT. This means that in human teeth the success rate determined by CBCT can be approximately 30% lower than that determined by periapical radiography. Paula-Silva *et al.* (2009a) endodontically treated 72 dog tooth roots and the outcome was evaluated at 6 months post-treatment. Unfavourable outcomes (emerged, unchanged or enlarged periapical lesions) were demonstrated in 21% of roots when analysed by periapical radiography, but in 65% when analysed by CBCT. This means that in dog teeth the percentage of favourable outcome determined by CBCT can be 40% lower than that determined by periapical radiography. These findings suggest that complete resolution of existing periapical radiolucency on radiographs does not guarantee a healthy periapex.

Paula-Silva *et al.* (2009b) evaluated the sensitivity, specificity, predictive values and accuracy of periapical radiography and CBCT in diagnosing apical periodontitis, using histopathologic findings as a gold standard. The negative predictive value (NPV) of periapical radiography in diagnosing apical periodontitis was 0.25; thus, 75% of cases confirmed healthy by periapical

radiography presented apical periodontitis by histology. The NPV of periapical radiography in diagnosing apical periodontitis was previously reported as 0.53 by Brynolf (1967), 0.55 by Rowe & Binnie (1974), 0.74 by Green *et al.* (1997), 0.20 by Ricucci & Langeland (1998), 0.67 by Barthel *et al.* (2004) and 0.35 by Stavropoulos & Wenzel (2007). The NPV of periapical radiography varied from 0.20 to 0.74 in four studies that analysed human materials (Brynolf 1967, Green *et al.* 1997, Ricucci & Langeland 1998, Barthel *et al.* 2004). The variability of NPV could be explained by different tooth positions of samples selected for different studies. In addition, different radiographic techniques were used (Brynolf 1967, Barthel *et al.* 2004). The NPV of CBCT in diagnosing apical periodontitis was almost twice as high as that of periapical radiography according to Paula-Silva *et al.* (2009b), indicating that CBCT more accurately detected a healthy periapex.

The incidence of periapical healing by scar after nonsurgical root canal treatment is low (Bhaskar 1966, Love & Firth 2009). Love & Firth (2009) performed apical surgery in 100 endodontically treated teeth with persistent periapical radiolucent lesions and found that the incidence of periapical granuloma, cyst, abscess and scar was 77%, 18%, 3% and 2% respectively. CBCT cannot be used to distinguish scar tissue from an inflammatory granuloma, therefore, one may question whether all CBCT detected radiolucencies are true lesions. In a study by Velvart *et al.* (2001), 78 CBCT-scanned human periapical lesions were confirmed to be true lesions during periapical surgery. In a study by Paula-Silva *et al.* (2009b), the positive predictive value was 1 for CBCT in diagnosing apical periodontitis in dogs teeth. This means that when a lesion was diagnosed by CBCT, 100% of the cases were periapically inflamed histologically.

Reduction of the size of radiolucency was considered to represent the healing of the periapical pathology and thus frequently used as radiographic criteria to determine successful treatment (Lewsey *et al.* 2001, Ng *et al.* 2007). Paula-Silva *et al.* (2009a) reported that amongst 30 cases where reduction in size of radiolucency was diagnosed by periapical radiography, 24 (80%) appeared as enlarged lesions in CBCT images. It was found that when lesions expanded in the cancellous bone, frequently in the lingual direction, the enlargement of the lesion could only be revealed by volumetric measurements using CBCT. The diagnosis of reduced periapical radiolucency with radiographs therefore does not guarantee that the healing process has begun or is continuing.

During recent years, progress was made in developing and improving digital radiography; with the new digital high resolution systems the detection of radiolucency may be improved. However, the improved image remains two-dimensional and subject to the limitations of anatomical noise.

Recall rates

The influence of low recall rates on the success rate was not addressed in most of the systematic reviews listed in Table 1.

The recall rate is defined as the percentage of patients that present for follow-up after treatment. Sixty-three clinical studies (1922–2002) were selected in a review by Ng *et al.* (2007), but only thirty-nine studies reported the recall rate. The median recall rate was 52.7%, and the lowest recall rate was 11% (Selden 1974).

According to Friedman (2002), 'When many subjects included in the inception cohort of a study are not available for follow-up, the unknown treatment outcome invalidates the results'. Sathorn *et al.* (2005) only reviewed studies where a high recall rate was reported.

Different terms were used for patients who were followed up after treatment (e.g., 'returnees'), whilst 'dropouts' and 'absentees' were the terms used to denote patients who were not present for follow-up. Ørstavik *et al.* (2004) compared the characteristics of returnees and dropout groups and found that dropout patients had more symptoms and perceived that treatment had failed. Therefore, in studies with large dropout groups, the reported success rate may represent an overestimation.

Efforts have been made to improve the recall rate. In the Toronto studies I, II and III (Friedman *et al.* 2003, Farzaneh *et al.* 2004b, Marquis *et al.* 2006), multiple letters and telephone calls were used to encourage patients to present themselves for follow-up; financial compensation was also offered for travel and loss of work time. In total, 1379 teeth were treated in these three studies; only 374 (27%) were included in the analysis. The outcome of the 73% of treated teeth is unknown; it therefore appears that the measures taken to increase the recall rate were not effective.

Ørstavik *et al.* (2004) suggested a new method to improve the overall recall rate. In that study, the percentage of dropouts increased from 33% at 1 year to 62% at 4 years. Over the 4-year period, 82% of patients were seen at least once, either at 1, 2, 3 or 4 years; only 18% of the patients never returned. Thus, the overall recall rate for the 4-year period reached 82%. However,

the year in which the outcome was recorded was not mentioned when patients were seen more than once.

Interestingly, the success rate usually increased over time, in parallel with a decreasing recall rate. The recall rate and success rate for each year during the follow-up period were provided by Ørstavik (1996) and are summarized in Table 2. In total, 155 teeth with preoperative periapical radiolucencies were treated and followed up for 4 years. The recall rate decreased from 71% at 1 year to 33% at 4 years. Meanwhile, the success rate increased from 44% at 1 year to 82% at 4 years. Thus, at 4 years, when only one out of three patients returned, the success rate was 82%. Such an increase in success rate over a 4-year period has been taken as evidence of a gradual periapical healing process (European Society of Endodontology 2006). However, another possible explanation could be that an increasing number of patients with unfavourable outcomes did not return for follow-up.

If this is true, then the 4-year follow-up period (European Society of Endodontology 2006) is unnecessary. Indeed, more than 88% of roots presenting reduced periapical radiolucency at 4 years following treatment had already shown this favourable outcome within 1 year (Ørstavik 1996). Wu & Wesselink (2005) suggested categorizing treatments where the existing radiolucent area was unchanged after 1 year as ineffective and requiring further treatment. Table 2 (Ørstavik 1996) shows that the recall rate at 1 year was more than twice as high as that at 4 years, suggesting that the overall recall rate could have been higher if the outcome had been determined at the end of 1 year. However, the suggestions made by Wu & Wesselink (2005) should be confirmed in studies utilizing CBCT.

Extractions and re-treatments

The systematic reviews listed in Table 1 did not discuss whether extractions and retreatments were included in the failure category.

Table 2 Success rate at 0–4 years post-treatment for 155 teeth with preoperative periapical radiolucency (Ørstavik 1996)

Year	Recall rate (%)	Success (PAI score of 1 or 2) rate (%)
0	100	0
1	71	44
2	60	72
3	55	79
4	33	82

During a period of 4 years or more between treatment and recall (European Society of Endodontology 2006, Ng *et al.* 2007), some teeth were extracted or retreated because of endodontic treatment-related complications (Molander *et al.* 1998, Cheung & Chan 2003). Petersson *et al.* (1991) found that 22% of teeth with failed root canal treatment were extracted. In a systematic review, Hepworth & Friedman (1997) found that 57% of retreatments were performed for 'technical' purposes, whilst 43% of retreatments were performed because of endodontic failures. Cheung & Chan (2003) categorized 314 root canal treated teeth as failures, including 143 (46%) extractions and 55 (18%) surgical or nonsurgical retreatments.

Zadik *et al.* (2008) analysed the reasons for extraction of root filled teeth. Endodontic treatment-related extractions comprised 30% of cases (iatrogenic perforation/stripping 9%, vertical root fracture 9%, endodontic failure 12%). All extractions were considered as failures by Ørstavik *et al.* (2004).

Amongst the 15 studies reviewed by Ng *et al.* (2007) in which the follow-up period was 4 years or longer (Cheung 2002, Chugal *et al.* 2001, Cvek 1992, Cvek *et al.* 1982, Grahnén & Hansen 1961, Halse & Molven 1987, Hoskinson *et al.* 2002, Reid *et al.* 1992, Smith *et al.* 1993, Ørstavik 1996, Sjögren *et al.* 1997, Werts 1975, Peak *et al.* 2001, Ørstavik *et al.* 1987, Sjögren *et al.* 1990), only one included extractions and retreatments in the failure category (Cheung 2002). In the other 14 studies, extractions were either not mentioned or excluded as the reason for extraction was unknown (Hoskinson *et al.* 2002); retreatments were not mentioned at all.

The PAI scoring system

In two reviews (Sathorn *et al.* 2005, Ng *et al.* 2007), the use of the periapical index (PAI) for the determination of success was described. However, the validity of using this index was not discussed in any of the systematic reviews listed in Table 1.

The PAI scoring system suggested by Ørstavik *et al.* (1986) has been used in 58 studies thus far (Table 3). Twenty of these studies were reported between 1987–2003. Thirty-eight were reported between 2004–2008. The number of such publications noticeably increased from one per year between 1987 and 2003, to seven per year between 2004 and 2008. The PAI scoring system seems therefore accepted as a valid tool to determine treatment outcome and to reveal changes in the extent and severity of periapical inflammation after root canal treatment.

Table 3 Fifty-eight studies using periapical index published between 1987 and 2008

Article
Ørstavik <i>et al.</i> 1987
Ørstavik <i>et al.</i> 2004
Ørstavik 1988
Ørstavik 1991
Ørstavik 1996
Eriksen <i>et al.</i> 1988a
Eriksen <i>et al.</i> 1988b
Ørstavik & Mjör 1992
Ørstavik & Hörsted-Bindslev 1993
Kerosuo & Ørstavik 1997
Valderhaug <i>et al.</i> 1997
Trope <i>et al.</i> 1999
Kirkevang <i>et al.</i> 2000
Kirkevang <i>et al.</i> 2001
Kirkevang <i>et al.</i> 2004
Kirkevang <i>et al.</i> 2006
Delano <i>et al.</i> 2001
Waltimo <i>et al.</i> 2001
Waltimo <i>et al.</i> 2005
Boucher <i>et al.</i> 2002
Dugas <i>et al.</i> 2003
Huunonen <i>et al.</i> 2003
Friedman <i>et al.</i> 2003
Kirkevang & Wenzel 2003
Camps <i>et al.</i> 2004
Jiménez-Pinzón <i>et al.</i> 2004
Farzaneh <i>et al.</i> 2004a
Farzaneh <i>et al.</i> 2004b
Peters <i>et al.</i> 2004
Segura-Egea <i>et al.</i> 2004
Segura-Egea <i>et al.</i> 2005
Segura-Egea <i>et al.</i> 2008
Wang <i>et al.</i> 2004
Yoldas <i>et al.</i> 2004
Chazel <i>et al.</i> 2005
Marending <i>et al.</i> 2005
Quesnell <i>et al.</i> 2005
Marquis <i>et al.</i> 2006
Peciuliene <i>et al.</i> 2006
Ridell <i>et al.</i> 2006
Skudutyte-Rysstad & Eriksen 2006
Terças <i>et al.</i> 2006
Conner <i>et al.</i> 2007
Frisk 2007
Ridao-Sacie <i>et al.</i> 2007
Simon <i>et al.</i> 2007
Cotton <i>et al.</i> 2008
de Chevigny <i>et al.</i> 2008a
de Chevigny <i>et al.</i> 2008b
Estrela <i>et al.</i> 2008a
Estrela <i>et al.</i> 2008b
Estrela <i>et al.</i> 2008c
Hannahan & Eleazer 2008
Holden <i>et al.</i> 2008
Kayahan <i>et al.</i> 2008
Penesis <i>et al.</i> 2008
Touré <i>et al.</i> 2008
Bahrami <i>et al.</i> 2008

In the PAI system, periapical radiolucencies are categorized with scores from 1 to 5: score 1 denotes a radiographically healthy periapex; scores 2–5 represent increasing extent and severity of apical periodontitis on radiography (Ørstavik *et al.* 1987). A relative incidence distribution (RIDIT, 0–1.0) is used to indicate the increasing severity of apical periodontitis at the histological level, with 0 representing absence of inflammation and 1.0 representing severe inflammation. According to findings from a histological and radiographical study of the periapical region of human maxillary incisors (Brynnolf 1967), the five scores for radiolucency correspond to the five different points of the RIDIT scale. For example, score 2 corresponds to 0.24 in the RIDIT (mild inflammation).

The PAI was based on only one study where the relationship between radiographic and histologic findings was reported (Brynnolf 1967). The thickness of the cortical bone and the distance from root tip to cortex varies remarkably for different tooth groups (Huumonen & Ørstavik 2002). The validity of applying the classification based on the maxillary incisor findings of Brynnolf (1967) to all tooth groups is thus questionable.

One of the reasons why PAI is gaining popularity could be that PAI allows the reduction in size of an existing lesion to be objectively determined. Ørstavik *et al.* (2004) reported that in 79% of roots, the PAI score decreased from a value of ≥ 3 preoperatively to a value of 1 or 2 post-treatment. Waltimo *et al.* (2001) reported that the mean RIDIT value decreased from 0.51 to 0.31 after treatment. However, the images on radiographs are two-dimensional, preventing the detection of bucco-lingual lesion expansion. Agbaje *et al.* (2007) and Paula-Silva *et al.* (2009a) used CBCT to measure the extraction sockets or periapical lesions; the volume was recorded in cubic millimetres. This represents a more reliable method for diagnosing reduced lesion size. Lesions frequently expanded in the cancellous bone and in lingual direction, such lesion enlargement was only revealed by volumetric measurements using CBCT (Paula-Silva *et al.* 2009a).

The rationale of including samples exhibiting a PAI score of 2 (corresponding to RIDIT 0.24) in the success category (Trope *et al.* 1999, Friedman *et al.* 2003, Farzaneh *et al.* 2004b, Ørstavik *et al.* 2004, Marquis *et al.* 2006) was probably the prevention of overdiagnosis of disease. However, considering that up to 53% of teeth had a score of 2 at the end of the study (Ørstavik *et al.* 2004), it is likely that many cases with small post-treatment lesions were included in the success category (Ørstavik *et al.* 1987). Ørstavik *et al.*

(2004) treated 192 roots with preoperative apical periodontitis. When PAI scores of 1 and 2 were considered to represent successful outcomes, the success rate was 79%; when only teeth exhibiting a PAI score of 1 were considered to represent successful outcomes, the success rate dropped to 26%.

Influencing factors

A number of factors that can influence treatment outcome were discussed in review articles; these factors included the presence of preoperative periapical radiolucency (Basmadjian-Charles *et al.* 2002, Ng *et al.* 2008a), root fillings with or without voids (Ng *et al.* 2008a), root fillings extending to 0–2 mm from the radiographic apex (Basmadjian-Charles *et al.* 2002, Kojima *et al.* 2004, Ng *et al.* 2008a), and the presence of satisfactory coronal restoration (Ng *et al.* 2008a). However, considering that the success rates reported in longitudinal studies might be overestimated, risk factor analysis should be performed again using CBCT to evaluate the data from new studies.

Concluding remarks

Several factors could have contributed to the overestimation of successful outcomes after root canal treatment:

1. A high percentage of cases that are confirmed healthy by periapical radiography present apical periodontitis in CBCT images and histology.
2. In teeth, where reduced size of the existing radiolucency is diagnosed by periapical radiography and considered to represent periapical healing, enlargement of the lesion may frequently be confirmed by CBCT.
3. PAI score 2 (mild inflammation) has usually been included in the success category.
4. Extractions and re-treatments were rarely recorded as failures.
5. The recall rate was often lower than 50% in longitudinal clinical studies.

Previously periapical radiography was the only imaging method available to diagnose post-treatment apical periodontitis. With the development of CBCT techniques, better understanding of the outcome of root canal treatment becomes possible. The outcomes of root canal treatment should be re-evaluated in long-term longitudinal studies using CBCT and stricter criteria. Systematic reviews repeatedly reporting overestimated success rates, as determined by periapical radiography

without considering the above mentioned limitations, mislead readers and could eventually damage the reputation of root canal treatment.

Important challenges remain: first, a consensus should be reached that the success rate reported in systematic reviews is likely to be an overestimation. However, longitudinal studies have shown that the periapical radiographs-determined success has a high prognostic validity (Fristad *et al.* 2004, Mead *et al.* 2005), which suggested that success may not require complete and utter radiographic resolution. A discussion on whether all asymptomatic post-treatment apical periodontitis should be treated and the replacement of periapical radiography with CBCT in clinics should follow. A more comprehensive understanding of the true effectiveness of current procedures will stimulate the development of new ideas and strategies, and thus improve the outcome and predictability of apical periodontitis treatment.

References

- Agbaje JO, Jacobs R, Maes F, Michiels K, van Steenberghe D (2007) Volumetric analysis of extraction sockets using cone beam computed tomography: a pilot study on ex vivo jaw bone. *Journal of Clinical Periodontology* **34**, 985–90.
- Bahrami G, Væth M, Kirkevang L-L, Wenzel A, Isidor F (2008) Risk factors for tooth loss in an adult population: a radiographic study. *Journal of Clinical Periodontology* **35**, 1059–65.
- Barthel CR, Zimmer S, Trope M (2004) Relationship of radiologic and histologic signs of inflammation in human root-filled teeth. *Journal of Endodontics* **30**, 75–9.
- Basmadjian-Charles CL, Farge P, Bourgeois DM, Lebrun T (2002) Factors influencing the long-term results of endodontic treatment: a review of the literature. *International Dental Journal* **52**, 81–6.
- Bender IB (1982) Factors influencing the radiographic appearance of bone lesions. *Journal of Endodontics* **8**, 161–70.
- Bender IB, Seltzer S (1961) Roentgenographic and direct observation of experimental lesions in bone: I. *Journal of American Dental Association* **62**, 152–60.
- Bhaskar SN (1966) Periapical lesions: types, incidence, and clinical features. *Oral Surgery, Oral Medicine, Oral Pathology* **21**, 657–71.
- Boucher Y, Matossian L, Rilliard F, Machtou P (2002) Radiographic evaluation of the prevalence and technical quality of root canal treatment in a French subpopulation. *International Endodontic Journal* **35**, 229–38.
- Brenner DJ, Hall EJ (2007) Computed tomography – an increasing source of radiation exposure. *The New England Journal of Medicine* **357**, 2277–84.
- Brynolf I (1967) A histological and roentgenological study of periapical region of human upper incisors. *Odontologisk Revy* **18**(Suppl. 11), 1–97.
- Camps J, Pommel L, Bukiet F (2004) Evaluation of periapical lesion healing by correction of gray values. *Journal of Endodontics* **30**, 762–6.
- Chazel JC, Tramini P, Valcarcel J, Pélissier B (2005) A comparative analysis of periapical health based on historic and current data. *International Endodontic Journal* **38**, 277–84.
- Cheung GS (2002) Survival of first-time nonsurgical root canal treatment performed in a dental teaching hospital. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **93**, 596–604.
- Cheung GSP, Chan TK (2003) Long-term survival of primary root canal treatment carried out in a dental teaching hospital. *International Endodontic Journal* **36**, 117–28.
- de Chevigny C, Dao TT, Basrani BR *et al.* (2008a) Treatment outcome in endodontics: the Toronto study—phases 3 and 4: orthograde retreatment. *Journal of Endodontics* **34**, 131–7.
- de Chevigny C, Dao TT, Basrani BR *et al.* (2008b) Treatment outcome in endodontics: the Toronto study—phase 4: initial treatment. *Journal of Endodontics* **34**, 258–63.
- Chugal NM, Clive JM, Spångberg LSW (2001) A prognostic model for assessment of the outcome of endodontic treatment: effect of biologic and diagnostic variables. *Oral Surgery, Oral Medicine, Oral Pathology, Radiology and Endodontics* **91**, 342–52.
- Conner DA, Caplan DJ, Teixeira FB, Trope M (2007) Clinical outcome of teeth treated endodontically with a nonstandardized protocol and root filled with resilon. *Journal of Endodontics* **33**, 1290–2.
- Cotton TP, Schindler WG, Schwartz SA, Watson WR, Hargreaves KM (2008) A retrospective study comparing clinical outcomes after obturation with Resilon/Epiphany or Gutta-Percha/Kerr sealer. *Journal of Endodontics* **34**, 789–97.
- Cvek M (1992) Prognosis of luxated non-vital maxillary incisors treated with calcium hydroxide and filled with gutta-percha. A retrospective clinical study. *Endodontics & Dental Traumatology* **8**, 45–55.
- Cvek M, Granath L, Lundberg M (1982) Failures and healing in endodontically treated non-vital anterior teeth with posttraumatically reduced pulpal lumen. *Acta Odontologica Scandinavica* **40**, 223–8.
- Delano EO, Ludlow JB, Ørstavik D, Tyndall D, Trope M (2001) Comparison between PAI and quantitative digital radiographic assessment of apical healing after endodontic treatment. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **92**, 108–15.
- Dugas NN, Lawrence HP, Teplitsky PE, Pharoah MJ, Friedman S (2003) Periapical health and treatment quality assessment of root-filled teeth in two Canadian populations. *International Endodontic Journal* **36**, 181–92.
- Eriksen HM, Bjertness E, Ørstavik D (1988a) Prevalence and quality of endodontic treatment in an urban adult population in Norway. *Endodontics & Dental Traumatology* **4**, 122–6.

- Eriksen HM, Ørstavik D, Kerekes K (1988b) Healing of apical periodontitis after endodontic treatment using three different root canal sealers. *Endodontics & Dental Traumatology* **4**, 114–7.
- Estrela C, Bueno MR, Leles CR, Azevedo B, Azevedo JR (2008a) Accuracy of cone beam computed tomography and panoramic and periapical radiography for detection of apical periodontitis. *Journal of Endodontics* **34**, 273–9.
- Estrela C, Bueno MR, Azevedo BC, Azevedo JR, Pécora JD (2008b) A new periapical index based on cone beam computed tomography. *Journal of Endodontics* **34**, 1325–31.
- Estrela C, Leles CR, Hollanda AC, Moura MS, Pécora JD (2008c) Prevalence and risk factors of apical periodontitis in endodontically treated teeth in a selected population of Brazilian adults. *Brazil Dental Journal* **19**, 34–9.
- European Society of Endodontology (2006) Quality guidelines for endodontic treatment: consensus report of the European Society of Endodontology. *International Endodontic Journal* **39**, 921–30.
- Farzaneh M, Abitbol S, Friedman S (2004a) Treatment outcome in endodontics: the Toronto study. Phases I and II: Orthograde retreatment. *Journal of Endodontics* **30**, 627–33.
- Farzaneh M, Abitbol S, Lawrence HP, Friedman S (2004b) Treatment outcome in endodontics—the Toronto Study. Phase II: initial treatment. *Journal of Endodontics* **30**, 302–9.
- Friedman S (2002) Considerations and concepts of case selection in the management of post-treatment endodontic disease (treatment failure). *Endodontic Topics* **1**, 54–78.
- Friedman S, Abitbol S, Lawrence HP (2003) Treatment outcome in endodontics: the Toronto Study. Phase I: initial treatment. *Journal of Endodontics* **29**, 787–93.
- Frisk F (2007) Epidemiological aspects on apical periodontitis. Studies based on the Prospective Population Study of Women in Göteborg and the Population Study on Oral Health in Jönköping. *Swedish Dental Journal Supplement* **189**, 11–78.
- Fristad I, Molven O, Halse A (2004) Nonsurgically retreated root-filled teeth – radiographic findings after 20–27 years. *International Endodontic Journal* **37**, 12–8.
- Grahnén H, Hansen L (1961) The prognosis of pulp and root canal therapy. *Odontologisk Revy* **12**, 146–65.
- Green TL, Walton RE, Taylor JK, Merrel P (1997) Radiographic and histologic periapical findings of root canal treated teeth in cadaver. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **83**, 707–11.
- Halse A, Molven O (1987) Overextended gutta-percha and Kloroperka N-Ø root canal fillings. Radiographic findings after 10–17 years. *Acta Odontologica Scandinavica* **45**, 171–7.
- Hannahan JP, Eleazer PD (2008) Comparison of success of implants versus endodontically treated teeth. *Journal of Endodontics* **34**, 1302–5.
- Hepworth MJ, Friedman S (1997) Treatment outcome of surgical and non-surgical management of endodontic failures. *Journal of Canadian Dental Association* **63**, 364–71.
- Holden DT, Schwartz SA, Kirkpatrick TC, Schindler WG (2008) Clinical outcomes of artificial root-end barriers with mineral trioxide aggregate in teeth with immature apices. *Journal of Endodontics* **34**, 812–7.
- Hoskinson SE, Ng YL, Hoskinson AE, Moles DR, Gulabivala K (2002) A retrospective comparison of outcome of root canal treatment using two different protocols. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **93**, 705–15.
- Huumonen S, Ørstavik D (2002) Radiological aspects of apical periodontitis. *Endodontic Topics* **1**, 3–25.
- Huumonen S, Lenander-Lumikari M, Sigurdsson A, Ørstavik D (2003) Healing of apical periodontitis after endodontic treatment: a comparison between a silicone-based and a zinc oxide-eugenol-based sealer. *International Endodontic Journal* **36**, 296–301.
- Jiménez-Pinzón A, Segura-Egea JJ, Poyato-Ferrera M, Velasco-Ortega E, Ríos-Santos JV (2004) Prevalence of apical periodontitis and frequency of root-filled teeth in an adult Spanish population. *International Endodontic Journal* **37**, 167–73.
- Jorge EG, Tanomaru-Filho M, Gonçalves M, Tanomaru JMG (2008) Detection of periapical lesion development by conventional radiography or computed tomography. *Oral Surgery Oral Medicine Oral Pathology Oral Radiology Endodontics* **106**, e56–61.
- Kayahan MB, Malkondu O, Canpolat C, Kaptan F, Bayirli G, Kazazoglu E (2008) Periapical health related to the type of coronal restorations and quality of root canal fillings in a Turkish subpopulation. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **105**, e58–62.
- Kerosuo E, Ørstavik D (1997) Application of computerised image analysis to monitoring endodontic therapy: reproducibility and comparison with visual assessment. *Dentomaxillofacial Radiology* **26**, 79–84.
- Kirkevang L-L, Wenzel A (2003) Risk indicators for apical periodontitis. *Community Dentistry and Oral Epidemiology* **31**, 59–67.
- Kirkevang L-L, Ørstavik D, Hörsted-Bindslev P, Wenzel A (2000) Periapical status and quality of root fillings and coronal restorations in a Danish population. *International Endodontic Journal* **33**, 509–15.
- Kirkevang L-L, Hörsted-Bindslev P, Ørstavik D, Wenzel A (2001) Frequency and distribution of endodontically treated teeth and apical periodontitis in an urban Danish population. *International Endodontic Journal* **34**, 198–205.
- Kirkevang LL, Vaeth M, Wenzel A (2004) Tooth-specific risk indicators for apical periodontitis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **97**, 739–44.
- Kirkevang LL, Vaeth M, Hörsted-Bindslev P, Wenzel A (2006) Longitudinal study of periapical and endodontic status in a Danish population. *International Endodontic Journal* **39**, 100–7.
- Kojima K, Inamoto K, Nagamatsu K et al. (2004) Success rate of endodontic treatment of teeth with vital and nonvital

- pulps. A meta-analysis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **97**, 95–9.
- Lewsey JD, Gilthorpe MS, Gulabivala K (2001) An introduction to meta-analysis within the framework of multilevel modeling using the probability of success of root canal treatment as an illustration. *Community Dental Health* **18**, 131–7.
- Lofthag-Hansen S, Huuononen S, Gröndahl K, Gröndahl HG (2007) Limited cone beam CT and intraoral radiography for the diagnosis of periapical pathology. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **103**, 114–9.
- Love RM, Firth N (2009) Histopathological profile of surgically removed persistent periapical radiolucent lesions of endodontic origin. *International Endodontic Journal* **42**, 198–202.
- Marending M, Peters OA, Zehnder M (2005) Factors affecting the outcome of orthograde root canal therapy in a general dentistry hospital practice. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **99**, 119–24.
- Marquis VL, Dao T, Farzaneh M, Abitbol S, Friedman S (2006) Treatment outcome in endodontics: the Toronto Study. Phase III: initial treatment. *Journal of Endodontics* **32**, 299–306.
- Mead C, Javidan-Nejad S, Mego ME, Nash B, Torabinejad M (2005) Levels of evidence for the outcome of endodontic surgery. *Journal of Endodontics* **31**, 19–24.
- Molander A, Reit C, Dahlén G, Kvist T (1998) Microbiological status of root-filled teeth with apical periodontitis. *International Endodontic Journal* **31**, 1–7.
- Ng Y-L, Mann V, Rahbaran S, Lewsey J, Gulabivala K (2007) Outcome of primary root canal treatment: systematic review of the literature – Part 1. Effects of study characteristics on probability of success. *International Endodontic Journal* **40**, 921–39.
- Ng Y-L, Mann V, Rahbaran S, Lewsey J, Gulabivala K (2008a) Outcome of primary root canal treatment: systematic review of the literature – Part 2. Influence of clinical factors. *International Endodontic Journal* **41**, 6–31.
- Ng Y-L, Mann V, Gulabivala K (2008b) Outcome of secondary root canal treatment: a systematic review of the literature. *International Endodontic Journal* **41**, 1026–46.
- Niederman R, Theodosopoulou N (2003) A systematic review of in vivo retrograde obturation materials. *International Endodontic Journal* **36**, 577–85.
- Ørstavik D (1988) Reliability of the periapical index scoring system. *Scandinavian Journal of Dental Research* **96**, 108–11.
- Ørstavik D (1991) Radiographic evaluation of apical periodontitis and endodontic treatment results: a computer approach. *International Dental Journal* **4**, 89–95.
- Ørstavik D (1996) Time-course and risk analyses of the development and healing of chronic apical periodontitis in man. *International Endodontic Journal* **29**, 150–5.
- Ørstavik D, Hörsted-Bindslev P (1993) A comparison of endodontic treatment results at two dental schools. *International Endodontic Journal* **26**, 348–54.
- Ørstavik D, Mjör IA (1992) Usage test of four endodontic sealers in *Macaca fascicularis* monkeys. *Oral Surgery, Oral Medicine, Oral Pathology* **73**, 337–44.
- Ørstavik D, Pitt Ford TR (1998) Apical periodontitis: microbial infection and host responses. In: Ørstavik D, Pitt Ford TR, eds. *Essential Endodontology*. Oxford: Blackwell Science, pp. 1–8.
- Ørstavik D, Kerekes K, Eriksen HM (1986) The periapical index: a scoring system for radiographic assessment of apical periodontitis. *Endodontics & Dental Traumatology* **2**, 20–34.
- Ørstavik D, Kerekes K, Eriksen HM (1987) Clinical performance of three endodontic sealers. *Endodontics & Dental Traumatology* **3**, 178–86.
- Ørstavik D, Qvist V, Stoltze K (2004) A multivariate analysis of the outcome of endodontic treatment. *European Journal of Oral Sciences* **112**, 224–30.
- Paik S, Sechrist C, Torabinejad M (2004) Levels of evidence for the outcome of endodontic retreatment. *Journal of Endodontics* **30**, 745–50.
- Peak JD, Hayes SJ, Bryant ST, Dummer PM (2001) The outcome of root canal treatment. A retrospective study within the armed forces (Royal Air Force). *British Dental Journal* **190**, 140–4.
- Peciuliene V, Rimkuvienė J, Maneliene R, Ivanauskaitė D (2006) Apical periodontitis in root filled teeth associated with the quality of root fillings. *Stomatologija* **8**, 122–6.
- Penesis VA, Fitzgerald PI, Fayad MI, Wenckus CS, BeGole EA, Johnson BR (2008) Outcome of one-visit and two-visit endodontic treatment of necrotic teeth with apical periodontitis: a randomized controlled trial with one-year evaluation. *Journal of Endodontics* **34**, 251–7.
- Peters OA, Barbakow F, Peters CI (2004) An analysis of endodontic treatment with three nickel–titanium rotary root canal preparation techniques. *International Endodontic Journal* **37**, 849–59.
- Peterson J, Gutmann JL (2001) The outcome of endodontic Resurgery: a systematic review. *International Endodontic Journal* **34**, 169–75.
- Petersson K, Håkansson R, Håkansson J, Olsson B, Wennberg A (1991) Follow-up study of endodontic status in an adult Swedish population. *Endodontics & Dental Traumatology* **7**, 221–5.
- Quesnell BT, Alves M, Hawkinson RW Jr, Johnson BR, Wenckus CS, BeGole EA (2005) The effect of human immunodeficiency virus on endodontic treatment outcome. *Journal of Endodontics* **31**, 633–6.
- Reid RJ, Abbott PV, McNamara JR, Heithersay GS (1992) A five-year study of hydron root canal fillings. *International Endodontic Journal* **25**, 213–20.
- Ricucci D, Bergenholtz G (2003) Bacterial status in root-filled teeth exposed to the oral environment by loss of restoration and fracture or caries – a histobacteriological study of treated cases. *International Endodontic Journal* **36**, 787–802.

- Ricucci D, Langeland K (1998) Apical limit of root canal instrumentation and obturation, part 2. A histological study. *International Endodontic Journal* **31**, 394–409.
- Ridao-Sacie C, Segura-Egea JJ, Fernández-Palacín A, Bullón-Fernández P, Ríos-Santos JV (2007) Radiological assessment of periapical status using the periapical index: comparison of periapical radiography and digital panoramic radiography. *International Endodontic Journal* **40**, 433–40.
- Ridell K, Petersson A, Matsson L, Mejäre I (2006) Periapical status and technical quality of root-filled teeth in Swedish adolescents and young adults. A retrospective study. *Acta Odontologica Scandinavica* **64**, 104–10.
- Rowe AHR, Binnie WH (1974) Correlation between radiological and histological changes following root canal treatment. *Journal of British Endodontic Society* **7**, 57–63.
- Sathorn C, Parashos P, Messer HH (2005) Effectiveness of single- versus multiple-visit endodontic treatment of teeth with apical periodontitis: a systematic review and meta-analysis. *International Endodontic Journal* **38**, 347–55.
- Segura-Egea JJ, Jiménez-Pinzón A, Poyato-Ferrera M, Velasco-Ortega E, Ríos-Santos JV (2004) Periapical status and quality of root fillings and coronal restorations in an adult Spanish population. *International Endodontic Journal* **37**, 525–30.
- Segura-Egea JJ, Jiménez-Pinzón A, Ríos-Santos JV, Velasco-Ortega E, Cisneros-Cabello R, Poyato-Ferrera M (2005) High prevalence of apical periodontitis amongst type 2 diabetic patients. *International Endodontic Journal* **38**, 564–9.
- Segura-Egea JJ, Jiménez-Pinzón A, Ríos-Santos JV, Velasco-Ortega E, Cisneros-Cabello R, Poyato-Ferrera MM (2008) High prevalence of apical periodontitis amongst smokers in a sample of Spanish adults. *International Endodontic Journal* **41**, 310–6.
- Selden HS (1974) Pulpoperiapical disease: diagnosis and healing. A clinical endodontic study. *Oral Surgery, Oral Medicine, Oral Pathology*, **27**, 271–83.
- Paula-Silva FWG, Hassan B, da Silva LAB, Leonardo MR, Wu M-K (2009a) Outcome of root canal treatment in dogs determined by periapical radiography and cone beam computed tomography. *Journal of Endodontics* **35**, 723–6.
- Paula-Silva FWG, Wu M-K, Leonardo MR, da Silva LAB, Wesselink PR (2009b) Accuracy of periapical radiography and cone beam computed tomography in diagnosing apical periodontitis using histopathological findings as a gold standard. *Journal of Endodontics*, in press
- Simon S, Rilliard F, Berdal A, Machtou P (2007) The use of mineral trioxide aggregate in one-visit apexification treatment: a prospective study. *International Endodontic Journal* **40**, 186–97.
- Sjögren U, Hägglund B, Sundqvist G, Wing K (1990) Factors affecting the long-term results of endodontic treatment. *Journal of Endodontics* **16**, 498–504.
- Sjögren U, Figdor D, Persson S, Sundqvist G (1997) Influence of infection at the time of root filling on the outcome of endodontic treatment of teeth with apical periodontitis. *International Endodontic Journal* **30**, 297–306.
- Skudutyte-Rysstad R, Eriksen HM (2006) Endodontic status amongst 35-year-old Oslo citizens and changes over a 30-year period. *International Endodontic Journal* **39**, 637–42.
- Smith CS, Setchell DJ, Hartly FJ (1993) Factors influencing the success of conventional root canal therapy – a five-year retrospective study. *International Endodontic Journal* **26**, 321–33.
- Stabholz A, Friedman S, Tamse A (1994) Endodontic failures and re-treatment. In: Cohen S, Burns RC, eds. *Pathways of the Pulp*, 6th edn. St Louis, MO: Mosby, pp. 692–3.
- Stavropoulos A, Wenzel A (2007) Accuracy of cone beam dental CT, intraoral digital and conventional film radiography for the detection of periapical lesion. An ex vivo study in pig jaws. *Clinical Oral Investigation* **11**, 101–6.
- van der Stelt PF (1985) Experimentally produced bone lesions. *Oral Surgery, Oral Medicine, Oral Pathology* **59**, 306–12.
- Tachibana H, Matsumoto K (1990) Applicability of X-ray computerized tomography in endodontics. *Endodontics & Dental Traumatology* **6**, 16–20.
- Terças AG, Oliveira AE, Lopes FF, Maia Filho EM (2006) Radiographic study of the prevalence of apical periodontitis and endodontic treatment in the adult population of São Luis, MA, Brazil. *Journal of Applied Oral Science* **14**, 183–7.
- Touré B, Kane AW, Sarr M, Ngom CT, Boucher Y (2008) Prevalence and technical quality of root fillings in Dakar, Senegal. *International Endodontic Journal* **41**, 41–9.
- Trope M (2003) The vital tooth-its importance in the study and practice of endodontics. *Endodontic Topics* **5**, 1–11.
- Trope M, Delano EO, Ørstavik D (1999) Endodontic treatment of teeth with apical periodontitis: single vs. multivisit treatment. *Journal of Endodontics* **25**, 345–50.
- Valderhaug J, Jokstad A, Ambjørnsen E, Norheim PW (1997) Assessment of the periapical and clinical status of crowned teeth over 25 years. *Journal of Dentistry* **25**, 97–105.
- Vandenbergh B, Jacobs R, Yang J (2008) Detection of periapical bone loss using digital intraoral and cone beam computed tomography images: an in vitro assessment of bony and/or infrabony defects. *Dentomaxillofacial Radiology* **37**, 252–60.
- Velvart P, Hecker H, Tillinger G (2001) Detection of the apical lesion and mandibular canal in conventional radiography and computed tomography. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **92**, 682–8.
- Waltimo TM, Boiesen J, Eriksen HM, Ørstavik D (2001) Clinical performance of 3 endodontic sealers. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **92**, 89–92.
- Waltimo T, Trope M, Haapasalo M, Ørstavik D (2005) Clinical efficacy of treatment procedures in endodontic infection control and one year follow-up of periapical healing. *Journal of Endodontics* **31**, 863–6.
- Wang N, Knight K, Dao T, Friedman S (2004) Treatment outcome in endodontics-The Toronto Study. Phases I and II: apical surgery. *Journal of Endodontics* **30**, 751–61.

- Werts R (1975) Endodontic treatment: a five-year follow-up. *Dental Survey* **51**, 29–30.
- Wu M-K, Wesslink PR (2005) Timeliness and effectiveness in the surgical management of persistent post-treatment periapical pathosis. *Endodontic Topics* **11**, 25–31.
- Wu M-K, Dummer PMH, Wesslink PR (2006) Consequences of and strategies to deal with residual post-treatment root canal infection. *International Endodontic Journal* **39**, 343–56.
- Yoldas O, Topuz A, Isçi AS, Oztunc H (2004) Postoperative pain after endodontic retreatment: single- versus two-visit treatment. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **98**, 483–7.
- Zadik Y, Sandler V, Bechor R, Salehrabi R (2008) Analysis of factors related to extractions of endodontically treated teeth. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology and Endodontics* **106**, e31–5.